arXiv:nlin/0102024v1 [nlin.SI] 20 Feb 2001

NTZ 1/2001

Universal $\mathbb{R}$-matrix as integral operator.

S.Derkachov[a,c1], D. Karakhanyan[b2] & R. Kirschner[c3]

[a] Department of Mathematics, St Petersburg Technology Institute,
Sankt Petersburg, Russia

[b] Yerevan Physics Institute ,
Br.Alikhanian st.2, 375036, Yerevan , Armenia.

[c] Naturwissenschaftlich-Theoretisches Zentrum und Institut für Theoretische Physik,
Universität Leipzig,
Augustusplatz 10, D-04109 Leipzig, Germany

Abstract.

We derive the integral operator form for the general rational solution of the Yang-Baxter equation with $s\ell(2|1)$ symmetry. Considering the defining relations for the kernel of the R-operator as a system of second order differential equations we observe remarkable reduction to a system of simple first order equations. The obtained kernel of R-operator has a very simple structure. To illustrate all this in the simplest situation we treat also the $s\ell(2)$ case.

---

[1] e-mail: Sergey.Derkachov@itp.uni-leipzig.de
[2] e-mail: karakhan@lx2.yerphi.am
[3] e-mail:Roland.Kirschner@itp.uni-leipzig.de

Contents

1 Introduction 2

2 $s\ell(2)$-invariant $\mathbb{R}$-matrix 4
  2.1 Algebra $s\ell(2)$ and lowest weight modules 4
  2.2 Yang-Baxter equation and general operator $\mathbb{R}_{\ell_1\ell_2}(u)$ 5
  2.3 Eigenvalues of operator $\mathbb{R}_{\ell_1\ell_2}(u)$ 6

3 The operator $\mathbb{R}_{\ell_1\ell_2}(u)$ as integral operator 7
  3.1 Differential equations 8
  3.2 The integral representation. 10

4 The supersymmetric extension: $s\ell(2|1)$-invariant $\mathbb{R}$-matrix 10
  4.1 The algebra $s\ell(2|1)$ and lowest weights modules 10
  4.2 Yang-Baxter equation and general operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$ 11
  4.3 Eigenvalues of operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$ 12

5 The operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$ as integral operator 13
  5.1 Differential equations 15
  5.2 Solution of the system of differential equations 17
  5.3 The integral representation. 19

6 Conclusions 20

7 Acknowledgments 20

1 Introduction

Solutions of the Yang-Baxter equation, in particular ones with $s\ell(2)$ or $s\ell(2|1)$ symmetry, are used in constructing a number of statistical models [1, 2, 3, 4, 5, 6, 8, 26] and are applied in analyzing the high-energy Regge [16, 17, 18, 22] and Bjorken asymptotics [19, 20, 21] of four -dimensional gauge theories.

In the previous paper [9] we have studied the $s\ell(2|1)$ symmetric rational solutions of the Yang-Baxter equation. We have obtained the general R-operator acting on the tensor product of two arbitrary representation spaces of lowest weight in terms of the matrix elements in a basis of lowest weight vectors appearing in the tensor product.

The lowest weight representation spaces have been chosen in terms of polynomials in one even variable z and two odd variables $\theta, \overline{\theta}$. Then the tensor product representation consists of polynomial two-point functions, i.e. functions of two sets $(z_i, \theta_i, \overline{\theta}_i)$. This formulation is the appropriate one for treating arbitrary infinite dimensional lowest weight representations. It is the natural formulation from the viewpoint of applications to the Bjorken limit of four-dimensional field theories like QCD. In the Bjorken asymtotics the interaction reduces in dimension effectively to a line on the light cone, being the range of the even variable z. The Grassmann variables $\theta, \overline{\theta}$ appear in the Bjorken limit of supersymmetric ($\mathcal{N} = 1$) Yang-Mills theories allowing to unify the gluon and quark fields in terms of superfields. The effective interaction can be formulated in terms of integral operators, the kernels of which are (super)conformal four-point functions.

This physical context provides a motivation to look for a corresponding formulation of the R-operators in terms of integral kernels. Instead of constructing these kernels from the matrix elements derived earlier we present here a direct calculation. We consider the defining relations for the general R-operator now as differential equations on the kernel. The essential observation is that this set of differential equation reduces to simple first order differential equations, one for each variable involved. The resulting kernel is expressed in powers of combinations of the variables having simple $s\ell(2|1)$ transformation properties.

The method proposed here seems to be general enough to become useful in solving the Yang-Baxter equation with other symmetries. In particular the observed reduction of the defining relation to simple first order differential equations is expected to work as a powerful tool in other cases. This reduction relies on the structure of the Lax operator, i.e. the R-operator acting in the tensor product of the fundamental and an arbitrary representation. Its matrix representation decomposes into matrices times derivatives, where these matrices are tensor products of particular vectors. In the case of $s\ell(2)$ this structure appeared earlier in calculations by L.N.Lipatov [16] and has been used in [17, 18, 22]. The kernel of R-operator is finally obtained as the similarity transformation transforming these first order differential operators into the simple derivatives with respect to variables $z_i, \theta_i, \bar{\theta}_i$.

Aiming to give a clear presentation of the method we discuss the simplest case of $s\ell(2)$ first. Note that the main idea to represent the universal R-matrix as an integral operator originates from the paper [10]. Moreover, the final answer for the kernel of R-operator has been obtained by a straightforward solution of the system of differential equations of the second order in the paper [11]. See this paper for the applications of the integral representation of R-matrix.

The presentation is organized as follows. In Section 2 we collect the standard facts about the algebra $s\ell(2)$ and its representations. We represent the lowest weight modules by polynomials in one variable (z) and the $s\ell(2)$-generators as first order differential operators. We derive the defining relation for the general R-matrix, i.e. the solution of the Yang-Baxter equation acting on tensor products of two arbitrary representations, the elements of which are polynomial functions of z_1 and z_2. We solve this defining relation in the space of lowest weights. The result is well known [28, 24, 25, 27] but we represent all calculations in the form which is appropriate for the supersymmetric generalisation. In Section 3 we give the simple method of derivation of the integral operator representation for the R-matrix. The main observation is that the defining system of differential equations of the second order for the kernel of R-matrix is equivalent to the system of the first-order differential equations.

These sections have introductory character and starting from the Section 4 we follow the same strategy for the supersymmetric algebra $s\ell(2|1)$. We represent the $s\ell(2|1)$ lowest weight modules by polynomials in one even (z) and two odd variables $(\theta, \bar{\theta})$ and the $s\ell(2|1)$-generators as first order differential operators and derive the defining relation for the general R-matrix, i.e. the solution of the Yang-Baxter equation acting on tensor products of two arbitrary representations, the elements of which are polynomial functions of $(z_1, \theta_1, \bar{\theta}_1)$ and $(z_2, \theta_2, \bar{\theta}_2)$. We solve this defining relation in the space of lowest weights. All this is the short summary of our previous paper [9] where the reader can find all details.

In Section 5 we reduce the defining system of differential equations of the second order for the kernel of R-matrix to the equivalent system of the first-order differential equations. Next we demonstrate by direct calculation that the obtained integral operator coincides with ones obtained in our previous paper.

Finally, in Section 6 we summarize. Appendix A contains some technical details about the construction of superconformal N-point functions. In Appendix B we give the list of superintegrals which are needed for the calculations of matrix elements.

2 $s\ell(2)$-invariant $\mathbb{R}$-matrix

2.1 Algebra $s\ell(2)$ and lowest weight modules

The algebra $s\ell(2)$ has three generators $\vec{S} = (S^1, S^2, S^3)$ with standard commutation relations:

$$[S^j, S^k] = i\epsilon^{jkp} S^p$$

We shall use the generators $S, S^{\pm}$:

$$S^{\pm} = S^1 \pm iS^2 \ , \ S = S^3 \ ; \ [S, S^{\pm}] = \pm S^{\pm} \ , \ [S^+, S^-] = 2S.$$

which have the following form in the fundamental representation

$$S^i = \frac{1}{2}\sigma^i \ ; \ \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \ ; \ \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \ ; \ \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$S^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \ ; \ S^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \ ; \ S = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

We represent the generators as first order differential operators, acting on the space of polynomials $\Phi(z)$:

$$S^- = -\partial \ ; \ S^+ = z^2\partial + 2\ell z \ ; \ S = z\partial + \ell. \tag{2.1.1}$$

Below we shall use the global form of $s\ell(2)$-transformations

$$e^{\lambda S^-}\Phi(z) = \Phi(z - \lambda) \ ; \ e^{\lambda S^+}\Phi(z) = \frac{1}{(1-\lambda z)^{2\ell}}\Phi\left(\frac{z}{1-\lambda z}\right) \tag{2.1.2}$$

The lowest weight $s\ell(2)$-module V_ℓ is built on the lowest weight vector ψ obeying:

$$S^-\psi = 0 \ ; \ S\psi = \ell\psi$$

The center of the enveloping algebra of the algebra $s\ell(2)$ is generated by the Casimir operator C_2 and the module is characterised uniquely by the action of the Casimir operators on its elements:

$$C_2 = S^2 - S + S^+S^- \ ; \ C_2 v = \ell(\ell - 1)v. \tag{2.1.3}$$

It is a vector space spanned by the following basis $V_k = (S^+)^k\psi \ , \ k \in \mathcal{Z}_+$. We shall use the above realization of the $s\ell(2)$-generators as the differential operators of first order acting on the infinite-dimensional(for generic ℓ) space V_ℓ of polynomials $\Phi(z)$ of variable z. The coherent state $e^{\lambda S^+}\psi$ for the lowest weight $\psi = 1$ is the generating function, the power expansion in λ of which produces the basis:

$$e^{\lambda S^+}1 = (1 - \lambda z)^{-2\ell} \ ; \ V_k = (2\ell)_k z^k \tag{2.1.4}$$

There exist some special values of ℓ: $\ell = -n \ ; \ n \in \frac{1}{2}\mathcal{Z}_+$, for which the module V_ℓ becomes a finite-dimensional vector space. Indeed it is evident from (2.1.4) that all basis vectors are equal zero for $k \geq n + 1$. Let us introduce the special notation for the fundamental $s\ell(2)$-module: $V_{-\frac{1}{2}} \equiv V_f$.

The tensor product of two $s\ell(2)$-modules has the well known direct sum decomposition:

$$V_{\ell_1} \otimes V_{\ell_2} = \sum_{n=0}^{\infty} V_{\ell_1+\ell_2+n} \tag{2.1.5}$$

For the proof of (2.1.5) in the generic situation one has to determine all possible lowest weight vectors appearing in the tensor product $V_{\ell_1}\otimes V_{\ell_2}$. In the realization on functions of z the space $V_{\ell_1}\otimes V_{\ell_2}$ is isomorphic to the space of polynomials in two variables $\psi(z_1,z_2)$. The $s\ell(2)$-generators acting on the $V_{\ell_1}\otimes V_{\ell_2}$ are just the sums of corresponding generators acting in V_{ℓ_i}. The lowest weight vectors of the irreducible representations in the decomposition of $V_{\ell_1}\otimes V_{\ell_2}$ are defined as the solutions of the equation:

$$S^-\Psi = 0 \Rightarrow \Psi(z_1,z_2) = \Psi\left(z_1-z_2\right) \tag{2.1.6}$$

Now, the lowest weight vectors in the decomposition of the tensor product are constructed from functions (2.1.6) being eigenfunctions of generator S. The eigenfunctions of the operator S are homogeneous polynomials of degree n , $n\in\mathcal{Z}_+$ and finally we obtain that all lowest weights in the space $V_{\ell_1}\otimes V_{\ell_2}$ are:

$$\Psi_n \equiv (z_1-z_2)^n \tag{2.1.7}$$

2.2 Yang-Baxter equation and general operator $\mathbb{R}_{\ell_1\ell_2}(u)$

Let V_{ℓ_i} ; $i=1,2,3$ be three lowest weight $s\ell(2)$-modules. Let us consider the three operators $\mathbb{R}_{\ell_i\ell_j}(u)$ which are acting in $V_{\ell_i}\otimes V_{\ell_j}$ and obey the Yang-Baxter equation in the space $V_{\ell_3}\otimes V_{\ell_1}\otimes V_{\ell_2}$:

$$\mathbb{R}_{\ell_3\ell_1}(u)\mathbb{R}_{\ell_3\ell_2}(v)\mathbb{R}_{\ell_1\ell_2}(v-u) = \mathbb{R}_{\ell_1\ell_2}(v-u)\mathbb{R}_{\ell_3\ell_2}(v)\mathbb{R}_{\ell_3\ell_1}(u) \tag{2.2.1}$$

To obtain the defining relation for the general $\mathbb{R}$-operator we consider the special case $\ell_1=f$ in (2.2.1)

$$\mathbb{R}_{f\ell_1}(u)\mathbb{R}_{f\ell_2}(v)\mathbb{R}_{\ell_1\ell_2}(v-u) = \mathbb{R}_{\ell_1\ell_2}(v-u)\mathbb{R}_{f\ell_2}(v)\mathbb{R}_{f\ell_1}(u). \tag{2.2.2}$$

We can choose the matrix realization in V_f and use the operator ($\vec{S}$ are generators acting in V_ℓ)

$$\mathbb{R}_{f\ell}(u) = u+\frac{1}{2}+\vec{\sigma}\vec{S} = \left(\begin{array}{cc} u+\frac{1}{2}+S & S^- \\ S^+ & u+\frac{1}{2}-S \end{array}\right), \tag{2.2.3}$$

which is the well known solution of the Yang-Baxter equation:

$$\mathbb{R}_{f_1f_2}(u-v)\mathbb{R}_{f_1\ell_3}(u)\mathbb{R}_{f_2\ell_3}(v) = \mathbb{R}_{f_2\ell_3}(v)\mathbb{R}_{f_1\ell_3}(u)\mathbb{R}_{f_1f_2}(u-v)\ ;\ \mathbb{R}_{f_1f_2}(u) = 1+u\cdot\mathbb{P}_{12},$$

where $\mathbb{P}_{12}$ is the permutation. The operators $\mathbb{R}_{f\ell_1}$,$\mathbb{R}_{f\ell_2}$ are linear functions of the spectral parameter u

$$\mathbb{R}_{f\ell_i}(u) = u+\frac{1}{2}+\mathbb{R}_i\ ;\ \mathbb{R}_i = \left(\begin{array}{cc} S_i & S_i^- \\ S_i^+ & -S_i \end{array}\right)\ ;\ i=2,3.$$

Now the general R-matrix $\mathbb{R}_{\ell_1\ell_2}(u)$ acting in the tensor product $V_{\ell_1}\otimes V_{\ell_2}$ of arbitrary modules, is fixed by the condition (2.2.2). After separation of $v+u$ and $v-u$ dependence we obtain two equations($v-u\rightarrow u$):

$$(\mathbb{R}_1+\mathbb{R}_2)\mathbb{R}_{\ell_1\ell_2}(u) = \mathbb{R}_{\ell_1\ell_2}(u)(\mathbb{R}_1+\mathbb{R}_2) \tag{2.2.4}$$

$$\left(\frac{u}{2}\left(\mathbb{R}_1-\mathbb{R}_2\right)+\mathbb{R}_1\mathbb{R}_2\right)\mathbb{R}_{\ell_1\ell_2}(u) = \mathbb{R}_{\ell_1\ell_2}(u)\left(\frac{u}{2}\left(\mathbb{R}_1-\mathbb{R}_2\right)+\mathbb{R}_2\mathbb{R}_1\right) \tag{2.2.5}$$

The first equation (2.2.4) expresses the fact that $\mathbb{R}_{\ell_1\ell_2}(u)$ has to be invariant with respect to the action of $s\ell(2)$-algebra

$$\left[\vec{S},\mathbb{R}_{\ell_1\ell_2}(u)\right] = 0\ ;\ \vec{S} = \vec{S}_1+\vec{S}_2$$

and the second equation is the wanted defining relation for the operator $\mathbb{R}_{\ell_1\ell_2}(u)$. In the matrix form the defining relation reads as follows:

$$K_{AB}\mathbb{R}_{\ell_1\ell_2}(u) = \mathbb{R}_{\ell_1\ell_2}(u)\bar{K}_{AB} \ ; \ A, B = 1, 2 \tag{2.2.6}$$

where:

$$K = \begin{pmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{pmatrix} \equiv \frac{u}{2}(\mathbb{R}_1 - \mathbb{R}_2) + \mathbb{R}_1\mathbb{R}_2 \ ; \ \bar{K} \equiv \frac{u}{2}(\mathbb{R}_1 - \mathbb{R}_2) + \mathbb{R}_2\mathbb{R}_1.$$

Due to $s\ell(2)$-invariance there is only one independent equation so that the system of equations (2.2.4), (2.2.5) for R-matrix is equivalent to the system

$$K_{12}\mathbb{R}_{\ell_1\ell_2}(u) = \mathbb{R}_{\ell_1\ell_2}(u)\bar{K}_{12} \ ; \ \left[\vec{S}, \mathbb{R}_{\ell_1\ell_2}(u)\right] = 0 \tag{2.2.7}$$

Indeed, the operators $\mathbb{R}_{f\ell_1}(u)\mathbb{R}_{f\ell_2}(v)$ and $\mathbb{R}_{f\ell_2}(v)\mathbb{R}_{f\ell_1}(u)$ are $s\ell(2)$-invariant by construction

$$\left[\vec{S} + \frac{1}{2}\vec{\sigma}, \mathbb{R}_{f\ell_1}(u)\mathbb{R}_{f\ell_2}(v)\right] = 0 \ ; \ \left[\vec{S} + \frac{1}{2}\vec{\sigma}, \mathbb{R}_{f\ell_2}(v)\mathbb{R}_{f\ell_1}(u)\right] = 0$$

so that we obtain the commutation relations between $s\ell(2)$-generators and K-operators

$$\left[\vec{S}, K\right] = -\frac{1}{2}\left[\vec{\sigma}, K\right] \ ; \ \left[\vec{S}, \bar{K}\right] = -\frac{1}{2}\left[\vec{\sigma}, \bar{K}\right].$$

For example

$$\left[S^+, K_{12}\right] = K_{11} - K_{22} \ , \ \left[S^+, K_{11}\right] = -K_{21} \ , \ \left[S^+, K_{22}\right] = K_{21} \ , \ \left[S^+, K_{21}\right] = 0$$

and it is possible to obtain the full system of equations (2.2.6) starting from the K_{12}-equation in (2.2.7).

2.3 Eigenvalues of operator $\mathbb{R}_{\ell_1\ell_2}(u)$

The $s\ell(2)$-invariance of the operator $\mathbb{R}_{\ell_1\ell_2}(u)$ allows to simplify the eigenvalue problem. Due to $s\ell(2)$-invariance any eigenspace of the operator $\mathbb{R}_{\ell_1\ell_2}$ is a lowest weight $s\ell(2)$-module generated by some lowest weight eigenvector. Therefore without loss of generality we can solve the defining relation (2.2.5) in the space of lowest weights. Let us consider in more details the structure of eigenspace of the $s\ell(2)$-invariant operator acting on the tensor product $V_{\ell_1} \otimes V_{\ell_2}$. As we have seen from the direct sum decomposition (2.1.5) for any fixed n the space of lowest weight vectors is one-dimensional and therefore the operator $\mathbb{R}_{\ell_1\ell_2}$ is diagonal on lowest weight vectors Ψ_n:

$$\mathbb{R}_{\ell_1\ell_2}(u)\Psi_n = R_n\Psi_n \tag{2.3.1}$$

The relation (2.2.7) leads to a simple recurrence relation for the eigenvalues R_n. The operator K_{12} commutes with the lowering generator S^- and therefore K_{12} maps lowest weight vectors to lowest weight vector and decrease its power by one:

$$K_{12}\Psi_n = \alpha_n(u)\Psi_{n-1} \ ; \ \alpha_n(u) = n(-u + \ell_1 + \ell_2 + n - 1),$$

where the concrete form of the function $\alpha_n(u)$ is derived by the explicit calculation using

$$K_{12} = -(z_1 - z_2)\partial_1\partial_2 - \left(\frac{u}{2} - \ell_2\right)\partial_1 + \left(\frac{u}{2} - \ell_1\right)\partial_2 \ ; \ \Psi_n = (z_1 - z_2)^n.$$

Under the permutation:

$$\ell_1, z_1 \leftrightarrow \ell_2, z_2 \ ; \ u \leftrightarrow -u,$$

the operators K_{AB} and lowest weights transform as follows:

$$K_{AB} \leftrightarrow \bar{K}_{AB} \ ; \ \Psi_n \leftrightarrow (-1)^{n+1}\Psi_n$$

so that the action of operator $\bar{K}_{12}$ on lowest weight vectors can be obtained from formulae for K_{12} by the formal substitution $\ell_1, z_1 \leftrightarrow \ell_2, z_2$ and $u \leftrightarrow -u$:

$$\bar{K}_{12}\Psi_n = -\alpha_n(-u)\Psi_{n-1}.$$

We project the operator equation $K_{12}\mathbb{R} = \mathbb{R}\bar{K}_{12}$ onto the lowest weight vectors Ψ_n:

$$K_{12}\mathbb{R}\Psi_n = \mathbb{R}\bar{K}_{12}\Psi_n \Longrightarrow \alpha_n(u)R_n\Psi_{n-1} = -R_{n-1}\alpha_n(-u)\Psi_{n-1}$$

which results in the recurrent relation:

$$\alpha_n(u)R_n = -\alpha_n(-u)R_{n-1}$$

with the general solution [28, 24, 25, 27]:

$$R_n(u) = (-1)^n \frac{\Gamma\left(-u+\ell_1+\ell_2\right)}{\Gamma\left(u+\ell_1+\ell_2\right)} \frac{\Gamma\left(u+\ell_1+\ell_2+n\right)}{\Gamma\left(-u+\ell_1+\ell_2+n\right)} \tag{2.3.2}$$

where we fixed the overall normalisation of R-operator in such a way that $\mathbb{R} : 1 \mapsto 1$.

3 The operator $\mathbb{R}_{\ell_1\ell_2}(u)$ as integral operator

It is useful to derive the explicit representation of the $\mathbb{R}$-matrix as integral operator acting on polynomials $\psi(z_1, z_2) \in V_{\ell_1} \otimes V_{\ell_2}$

$$\left[\mathbb{R}_{\ell_1\ell_2}(u)\psi\right](z_1, z_2) = \int\int \mathrm{d}z_3\mathrm{d}z_4 \mathcal{R}(z_1, z_2|z_3, z_4)\psi(z_3, z_4) \tag{3.0.1}$$

We suppose that it is possible to integrate by parts so that the equations (2.2.4), (2.2.5) result in the set of differential equations for the kernel $\mathcal{R}(\vec{z}) = \mathcal{R}(z_1, z_2|z_3, z_4)$

$$(\mathbb{R}_1 + \mathbb{R}_2 + \mathbb{R}_3 + \mathbb{R}_4)\mathcal{R}(\vec{z}) = 0 \tag{3.0.2}$$

$$\left(\frac{u}{2}\left(\mathbb{R}_1 - \mathbb{R}_2\right) + \mathbb{R}_1\mathbb{R}_2\right)\mathcal{R}(\vec{z}) = \left(\frac{u}{2}\left(\mathbb{R}_4 - \mathbb{R}_3\right) + \mathbb{R}_4\mathbb{R}_3\right)\mathcal{R}(\vec{z}) \tag{3.0.3}$$

where the matrices $\mathbb{R}_i \ , \ i = 3, 4$ arise after integration by parts

$$\mathbb{R}_i = \begin{pmatrix} S_i & S_i^- \\ S_i^+ & -S_i \end{pmatrix} = \begin{pmatrix} z_i\partial_i + \ell_i & -\partial_i \\ z_i^2\partial_i + 2\ell_i z_i & -z_i\partial_i - \ell_i \end{pmatrix} \ ; \ \ell_3 \equiv 1 - \ell_1 \ , \ \ell_4 \equiv 1 - \ell_2$$

First we shall solve this system of differential equations and then specify selfconsistently the contours of integration. The main system of equations can be simplified drastically.

Proposition 1 *The system of differential equations (3.0.2), (3.0.3) of the second order is equivalent to the system of differential equations of the first order*

$$\mathcal{D}_1\mathcal{R} = 0 \ , \ \mathcal{D}_2\mathcal{R} = 0 \ , \ \mathcal{D}_3\mathcal{R} = 0 \ , \ \mathcal{D}_4\mathcal{R} = 0. \tag{3.0.4}$$

where operators $\mathcal{D}_i$ are defined in a following way

$$\mathcal{D}_1 \equiv \partial_1 + \frac{u+\ell_1+\ell_4}{z_{14}} - \frac{u+\ell_4-\ell_1}{z_{12}} \ ; \ \mathcal{D}_2 \equiv \partial_2 + \frac{u+\ell_2+\ell_3}{z_{23}} + \frac{u+\ell_3-\ell_2}{z_{12}} \tag{3.0.5}$$

$$\mathcal{D}_3 \equiv \partial_3 - \frac{u+\ell_2+\ell_3}{z_{23}} - \frac{u+\ell_2-\ell_3}{z_{34}} \ ; \ \mathcal{D}_4 \equiv \partial_4 - \frac{u+\ell_1+\ell_4}{z_{14}} + \frac{u+\ell_1-\ell_4}{z_{34}}.$$

The $s\ell(2)$-generators are expressed in terms of $\mathcal{D}_i$ as follows:

$$\mathbb{S}^- = -\sum_{i=1}^4 \mathcal{D}_i \ ; \ \mathbb{S} = \sum_{i=1}^4 z_i\mathcal{D}_i \ ; \ \mathbb{S}^+ = \sum_{i=1}^4 z_i^2\mathcal{D}_i \ ; \ \vec{\mathbb{S}} = \sum_{i=1}^4 \vec{S}_i$$

The common solution of the system (3.0.4) up to overall normalisation is

$$\mathcal{R}(\vec{z}) \sim z_{12}^{u-\ell_1+\ell_4} z_{14}^{-u-\ell_1-\ell_4} z_{32}^{-u-\ell_2-\ell_3} z_{34}^{u+\ell_2-\ell_3} \ ; \ \ell_3 = 1-\ell_1 \ , \ \ell_4 = 1-\ell_2.$$

3.1 Differential equations

There exists the simple and elegant way to derive the equation of the first order for the R-matrix which is especially useful for the supersymmetric generalisation. The similar trick was used in [16, 17, 18, 22] for the derivation of integrals of motion for XXX spin chains. The matrix $\mathbb{R}$ has very special structure

$$\mathbb{R} = \begin{pmatrix} z & -1 \\ z^2 & -z \end{pmatrix}\partial + \begin{pmatrix} \ell & 0 \\ 2\ell z & -\ell \end{pmatrix} \ ; \ \begin{pmatrix} z & -1 \\ z^2 & -z \end{pmatrix} = \begin{pmatrix} 1 \\ z \end{pmatrix}\begin{pmatrix} z & -1 \end{pmatrix}$$

so that the matrix

$$\mathbb{R}f(z) = \begin{pmatrix} z & -1 \\ z^2 & -z \end{pmatrix} f'(z) + \begin{pmatrix} \ell & 0 \\ 2\ell z & -\ell \end{pmatrix} f(z)$$

has two evident eigenvectors

$$\begin{pmatrix} z & -1 \end{pmatrix}\mathbb{R}f(z) = -\ell\begin{pmatrix} z & -1 \end{pmatrix} f(z) \ ; \ \mathbb{R}f(z)\begin{pmatrix} 1 \\ z \end{pmatrix} = \ell\begin{pmatrix} 1 \\ z \end{pmatrix} f(z)$$

where $f(z)$ is an arbitrary function. The derivative $f'(z)$ disappears from the eigenvalues and we shall use this property to derive the first order differential equation for the kernel $\mathcal{R}(\vec{z})$. Let us multiply the equations (2.2.4), (2.2.5) to the vector $\begin{pmatrix} z_4 & -1 \end{pmatrix}$ from the left and to the vector $\begin{pmatrix} 1 \\ z_2 \end{pmatrix}$ from the right. Then we obtain the system of two first-order differential equations for the function $\mathcal{R}(\vec{z})$

$$\mathbb{D}_1\mathcal{R} + \mathbb{D}_3\mathcal{R} = (\ell_2-\ell_4)z_{24}\mathcal{R} \ ; \ \left(\frac{u}{2}+\ell_2\right)\mathbb{D}_1\mathcal{R} + \left(\frac{u}{2}+\ell_4\right)\mathbb{D}_3\mathcal{R} = -\frac{u}{2}(\ell_2-\ell_4)z_{24}\mathcal{R}$$

which is equivalent to the system

$$\mathbb{D}_1\mathcal{R} = -\left(u+\ell_4\right)z_{24}\mathcal{R} \ ; \ \mathbb{D}_3\mathcal{R} = \left(u+\ell_2\right)z_{24}\mathcal{R}. \tag{3.1.6}$$

We use the notation

$$\mathbb{D}_k \equiv \begin{pmatrix} z_4 & -1 \end{pmatrix} \mathbb{R}_k \begin{pmatrix} 1 \\ z_2 \end{pmatrix} = -z_{k2} z_{k4} \partial_k - \ell_k (z_{k2} + z_{k4}) \ ; \ k = 1, 3$$

for the arising differential operators. Let us multiply the equations (2.2.4), (2.2.5) to the vector $\begin{pmatrix} z_1 & -1 \end{pmatrix}$ from the left and to the vector $\begin{pmatrix} 1 \\ z_3 \end{pmatrix}$ from the right. In the same manner we obtain the mirror system of equations:

$$\mathbb{D}_2 \mathcal{R} = -\left(u + \ell_3\right) z_{13} \mathcal{R} \ ; \ \mathbb{D}_4 \mathcal{R} = \left(u + \ell_1\right) z_{13} \mathcal{R}, \tag{3.1.7}$$

where

$$\mathbb{D}_k \equiv \begin{pmatrix} z_1 & -1 \end{pmatrix} \mathbb{R}_k \begin{pmatrix} 1 \\ z_3 \end{pmatrix} = -z_{k1} z_{k3} \partial_k - \ell_k (z_{k1} + z_{k3}) \ ; \ k = 2, 4.$$

Finally, as consequence of the main system of differential equations of the second order we obtain the system of differential equations (3.1.6), (3.1.7) of the first order. The eqs. (3.1.6), (3.1.7) are equivalent to

$$\mathcal{D}_1 \mathcal{R} = 0 \ , \ \mathcal{D}_2 \mathcal{R} = 0 \ , \ \mathcal{D}_3 \mathcal{R} = 0 \ , \ \mathcal{D}_4 \mathcal{R} = 0.$$

where the differential operators $\mathcal{D}_i$ are defined in (3.0.5).
The next step is to prove the reverse, that the system of equations (3.0.2), (3.0.3) is the consequence of the system (3.0.4). It is easy to rewrite the $s\ell(2)$-generators and the equations of $s\ell(2)$-invariance (3.0.2) in terms of the introduced operators $\mathcal{D}_i$:

$$\mathbb{S}^- = \sum_{i=1}^{4} S_i^- = -\sum_{i=1}^{4} \mathcal{D}_i \ ; \ \mathbb{S} = \sum_{i=1}^{4} S_i^- = \sum_{i=1}^{4} z_i \mathcal{D}_i \ ; \ \mathbb{S}^+ = \sum_{i=1}^{4} S_i^+ = \sum_{i=1}^{4} z_i^2 \mathcal{D}_i.$$

$$\mathbb{S}^- \mathcal{R} = \sum_{i=1}^{4} \mathcal{D}_i \mathcal{R} = 0 \ ; \ \mathbb{S} \mathcal{R} = \sum_{i=1}^{4} z_i \mathcal{D}_i \mathcal{R} = 0 \ ; \ \mathbb{S}^+ \mathcal{R} = \sum_{i=1}^{4} z_i^2 \mathcal{D}_i = 0$$

As explained above due to $s\ell(2)$-invariance we can restrict to one K_{12}-equation in (2.2.7). This equation has the explicit form

$$z_{12} \partial_1 \partial_2 \mathcal{R}(\vec{z}) + z_{34} \partial_3 \partial_4 \mathcal{R}(\vec{z}) =$$
$$= -\left(\frac{u}{2} - \ell_2\right) \partial_1 \mathcal{R}(\vec{z}) + \left(\frac{u}{2} - \ell_1\right) \partial_2 \mathcal{R}(\vec{z}) - \left(\frac{u}{2} - \ell_4\right) \partial_3 \mathcal{R}(\vec{z}) + \left(\frac{u}{2} - \ell_3\right) \partial_4 \mathcal{R}(\vec{z})$$

and can be easily rewritten in terms of the operators $\mathcal{D}_i$

$$z_{12} \mathcal{D}_1 \mathcal{D}_2 \mathcal{R} + z_{34} \mathcal{D}_3 \mathcal{D}_4 \mathcal{R} =$$
$$= \left(\frac{u}{2} + \ell_3 + (u + \ell_2 + \ell_3) \frac{z_{12}}{z_{23}}\right) \mathcal{D}_1 \mathcal{R} - \left(\frac{u}{2} + \ell_4 - (u + \ell_1 + \ell_4) \frac{z_{12}}{z_{14}}\right) \mathcal{D}_2 \mathcal{R} +$$
$$+ \left(\frac{u}{2} + \ell_1 - (u + \ell_1 + \ell_4) \frac{z_{34}}{z_{14}}\right) \mathcal{D}_3 \mathcal{R} - \left(\frac{u}{2} + \ell_2 + (u + \ell_2 + \ell_3) \frac{z_{34}}{z_{23}}\right) \mathcal{D}_4 \mathcal{R}.$$

We see that the system of equations (3.0.2), (3.0.3) is the consequence of the system (3.0.4) and finally, the system of the defining equation (2.2.7) is equivalent to the system (3.0.4).
The common solution of the system (3.0.4) up to an overall normalisation is

$$\mathcal{R}(\vec{z}) \sim z_{12}^{u - \ell_1 + \ell_4} z_{14}^{-u - \ell_1 - \ell_4} z_{32}^{-u - \ell_2 - \ell_3} z_{34}^{u + \ell_2 - \ell_3} \ ; \ \ell_3 = 1 - \ell_1 \ , \ \ell_4 = 1 - \ell_2.$$

The operators $\mathcal{D}_i$ are obtained by the simple similarity transformation from the derivatives ∂_i

$$\mathcal{D}_1 = z_{12}^{u+\ell_4-\ell_1} z_{14}^{-u-\ell_1-\ell_4} \partial_1 z_{12}^{-u-\ell_4+\ell_1} z_{14}^{u+\ell_1+\ell_4} \ , \ \mathcal{D}_2 = z_{21}^{u+\ell_3-\ell_2} z_{23}^{-u-\ell_2-\ell_3} \partial_2 z_{21}^{-u-\ell_3+\ell_2} z_{23}^{u+\ell_2+\ell_3}$$

$$\mathcal{D}_3 = z_{32}^{-u-\ell_2-\ell_3} z_{34}^{u+\ell_2-\ell_3} \partial_3 z_{32}^{u+\ell_2+\ell_3} z_{34}^{-u-\ell_2+\ell_3} \ , \ \mathcal{D}_4 = z_{41}^{-u-\ell_1-\ell_4} z_{43}^{u+\ell_1-\ell_4} \partial_4 z_{41}^{u+\ell_1+\ell_4} z_{43}^{-u-\ell_1+\ell_4} .$$

so that by using the ansatz

$$\mathcal{R}(\vec{z}) \sim z_{12}^{u-\ell_1+\ell_4} z_{14}^{-u-\ell_1-\ell_4} z_{32}^{-u-\ell_2-\ell_3} z_{34}^{u+\ell_2-\ell_3} \cdot R(\vec{z})$$

the system of equations $\mathcal{D}_i \mathcal{R}(\vec{z}) = 0$ can be reduced to the system $\partial_i R(\vec{z}) = 0$ with the evident solution $R(\vec{z}) = const.$

3.2 The integral representation.

It remains to fix the contours of integrations. The $\mathbb{R}$-operator acts on the space of polynomials $\psi(z_1, z_2)$ which can be of arbitrary large degree. The requirement of convergence allows compact contours only and there remains the one possibility:

$$[\mathbb{R}_{\ell_1 \ell_2}(u)\psi]\,(z_1, z_2) = \rho \int_{z_1}^{z_2} \mathrm{d}z_3 \int_{z_1}^{z_3} \mathrm{d}z_4 z_{21}^{1+u-\ell_1-\ell_2} z_{41}^{\ell_2-\ell_1-u-1} z_{23}^{\ell_1-\ell_2-u-1} z_{34}^{\ell_1+\ell_2+u-1} \psi(z_3, z_4) \tag{3.2.8}$$

where the normalisation factor ρ is chosen so that $\mathbb{R} : 1 \mapsto 1$,

$$\rho = \frac{1}{\Gamma(\ell_1 - \ell_2 - u)\Gamma(\ell_2 - \ell_1 - u)} \frac{\Gamma(\ell_1 + \ell_2 - u)}{\Gamma(\ell_1 + \ell_2 + u)}. \tag{3.2.9}$$

It is easy to check *a posteriori* that the obtained integral operator is $s\ell(2)$-invariant and has the eigenvalues (2.3.2). The eigenvalues can be calculated by using twice the general formula

$$\int_{z_1}^{z_3} \mathrm{d}z_4 z_{34}^{a-1} z_{41}^{b-1} = z_{31}^{a+b-1} \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

4 The supersymmetric extension: $s\ell(2|1)$-invariant $\mathbb{R}$-matrix

4.1 The algebra $s\ell(2|1)$ and lowest weights modules

Here we collect the main formulas about algebra $s\ell(2|1)$ (for details see [12, 13, 14, 15, 9]). We represent the generators as first order differential operators, acting on the space of polynomials $\Phi(z, \theta, \bar{\theta})$:

$$S^- = -\partial \ ; \ V^- = \partial_\theta + \frac{1}{2}\bar{\theta}\partial \ ; \ W^- = \partial_{\bar{\theta}} + \frac{1}{2}\theta\partial, \tag{4.1.1}$$

$$V^+ = -\left[z\partial_\theta + \frac{1}{2}\bar{\theta}z\partial + \frac{1}{2}\bar{\theta}\theta\partial_\theta\right] - (\ell - b)\bar{\theta} \ ; \ W^+ = -\left[z\partial_{\bar{\theta}} + \frac{1}{2}\theta z\partial + \frac{1}{2}\theta\bar{\theta}\partial_{\bar{\theta}}\right] - (\ell + b)\theta,$$

$$S^+ = z^2\partial + z\theta\partial_\theta + z\bar{\theta}\partial_{\bar{\theta}} + 2\ell z - b\theta\bar{\theta}, \tag{4.1.2}$$

$$S = z\partial + \frac{1}{2}\theta\partial_\theta + \frac{1}{2}\bar{\theta}\partial_{\bar{\theta}} + \ell \ ; \ B = \frac{1}{2}\bar{\theta}\partial_{\bar{\theta}} - \frac{1}{2}\theta\partial_\theta + b.$$

The lowest weight $s\ell(2|1)$-module $V_{\ell,b} = V_{\vec{\ell}}$, $\vec{\ell} = (\ell, b)$ is built on the lowest weight vector ψ obeying:

$$V_-\psi = 0 \ ; \ W^-\psi = 0 \ ; \ S^-\psi = 0 \ ; \ S\psi = \ell\psi \ ; \ B\psi = b\psi$$

The module $V_{\vec{\ell}}$ is a vector space spanned by the following basis [12], [14] with the even vectors

$$A_k = (S^+)^k\psi \ ; \ B_k = (S^+)^{k-1}W^+V^+\psi \ , \ k \in \mathcal{Z}_+$$

and the odd vectors

$$V_k = (S^+)^k V^+\psi \ ; \ W_k = (S^+)^k W^+\psi \ , \ k \in \mathcal{Z}_+$$

We shall use the above realization of the $s\ell(2|1)$-generators as the differential operators of first order acting on the infinite-dimensional(for generic ℓ) space $V_{\ell,b}$ of polynomials $\Phi(z,\theta,\bar{\theta})$ of variables $z,\theta,\bar{\theta}$.

The tensor product of two $s\ell(2|1)$-modules has the following direct sum decomposition [14, 9]:

$$V_{\ell_1,b_1}\otimes V_{\ell_2,b_2} = V_{\ell,b} + 2\sum_{n=1}^{\infty} V_{\ell+n,b} + \sum_{n=0}^{\infty} V_{\ell+n+\frac{1}{2},b-\frac{1}{2}} + \sum_{n=0}^{\infty} V_{\ell+n+\frac{1}{2},b+\frac{1}{2}} \ ; \ \ell_i \neq \pm b_i \tag{4.1.3}$$

$$\ell = \ell_1 + \ell_2 \ ; \ b = b_1 + b_2$$

Note that this formula is applicable in the generic situation $\ell_i \neq \pm b_i$. All possible lowest weight vectors appearing in the tensor product $V_{\ell_1,b_1}\otimes V_{\ell_2,b_2}$ are divided in two sets, the even lowest weights:

$$\Phi_n^{\pm} \equiv \left(Z_{12} \pm \frac{1}{2}\theta_{12}\bar{\theta}_{12}\right)^n \ ; \ D_1^{\pm}\Phi_n^{\pm} = 0 \ , \ S\Phi_n^{\pm} = (n+\ell)\Phi_n^{\pm} \ , \ B\Phi_n^{\pm} = b\Phi_n^{\pm} \tag{4.1.4}$$

and the odd lowest weights:

$$\Psi_n^- \equiv \theta_{12}Z_{12}^n \ ; \ \Psi_n^+ \equiv \bar{\theta}_{12}Z_{12}^n \ ; \ S\Psi_n^{\pm} = (n+\ell+\frac{1}{2})\Psi_n^{\pm} \ , \ B\Psi_n^{\pm} = (b \pm \frac{1}{2})\Psi_n^{\pm} \tag{4.1.5}$$

where

$$Z_{ij} \equiv z_i - z_j + \frac{1}{2}(\bar{\theta}_i\theta_j + \theta_i\bar{\theta}_j) \ ; \ \theta_{ij} \equiv \theta_i - \theta_j \ ; \ \bar{\theta}_{ij} \equiv \bar{\theta}_i - \bar{\theta}_j.$$

4.2 Yang-Baxter equation and general operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$

Let $V_{\ell_i,b_i} \ ; \ i = 1,2,3$ be three lowest weight $s\ell(2|1)$-modules. We shall use the short-hand notation:

$$\vec{f} = \left(-\frac{1}{2}, -\frac{1}{2}\right) \ ; \ \vec{\ell} = (\ell, b) \ ; \ V_{\vec{\ell}} = V_{\ell,b}.$$

Let us consider the three operators $\mathbb{R}_{\vec{\ell}_i\vec{\ell}_j}(u)$ which are acting in $V_{\vec{\ell}_i}\otimes V_{\vec{\ell}_j}$ and obey the Yang-Baxter equation in the space $V_{\vec{\ell}_3}\otimes V_{\vec{\ell}_1}\otimes V_{\vec{\ell}_2}$ [25]:

$$\mathbb{R}_{\vec{\ell}_3\vec{\ell}_1}(u)\mathbb{R}_{\vec{\ell}_3\vec{\ell}_2}(v)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(v-u) = \mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(v-u)\mathbb{R}_{\vec{\ell}_3\vec{\ell}_2}(v)\mathbb{R}_{\vec{\ell}_3\vec{\ell}_1}(u) \tag{4.2.1}$$

To obtain the defining relation for the general $\mathbb{R}$-operator we consider the special case $\vec{\ell}_3 = \vec{f}$ in (4.2.1). Then one can choose the matrix realization [26, 7, 9] in $V_{\vec{\ell}_1}$

$$\mathbb{R}_{\vec{f}\vec{\ell}}(u) = \begin{pmatrix} u+S+B & -W^- & S^- \\ V^+ & u+2B & V^- \\ S^+ & W^+ & u+B-S \end{pmatrix}$$

and the operators $\mathbb{R}_{f\vec{\ell}_1}$,$\mathbb{R}_{f\vec{\ell}_2}$ are linear functions of spectral parameter u in this particular case

$$\mathbb{R}_{f\vec{\ell}_i}(u) = u + \mathbb{R}_i \ ; \ \mathbb{R}_i = \begin{pmatrix} S_i + B_i & -W_i^- & S_i^- \\ V_i^+ & 2B_i & V_i^- \\ S_i^+ & W_i^+ & B_i - S_i \end{pmatrix} \ ; \ i = 1, 2$$

Now the general R-matrix $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$ acting in the tensor product $V_{\vec{\ell}_1} \otimes V_{\vec{\ell}_2}$ of arbitrary modules, is fixed by the condition

$$\mathbb{R}_{f\vec{\ell}_1}(u)\mathbb{R}_{f\vec{\ell}_2}(v)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(v-u) = \mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(v-u)\mathbb{R}_{f\vec{\ell}_2}(v)\mathbb{R}_{f\vec{\ell}_1}(u) \tag{4.2.2}$$

or equivalently:

$$(\mathbb{R}_1 + \mathbb{R}_2)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u) = \mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)(\mathbb{R}_1 + \mathbb{R}_2) \tag{4.2.3}$$

$$\left(\frac{u}{2}(\mathbb{R}_1 - \mathbb{R}_2) + \mathbb{R}_1\mathbb{R}_2\right)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u) = \mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)\left(\frac{u}{2}(\mathbb{R}_1 - \mathbb{R}_2) + \mathbb{R}_2\mathbb{R}_1\right) \tag{4.2.4}$$

The first equation (2.2.4) expresses the fact that $\mathbb{R}(u)$ has to be invariant with respect to the action of $s\ell(2|1)$-algebra and the second equation is the defining relation for the operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$.

4.3 Eigenvalues of operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$

Due to $s\ell(2|1)$-invariance any eigenspace of the operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}$ is a lowest weight $s\ell(2|1)$-module generated by some lowest weight eigenvector. As we have seen from direct sum decomposition (4.1.3) for every fixed n the space of lowest weight vectors with eigenvalue b is two-dimensional and the ones with eigenvalues $b \pm \frac{1}{2}$ are one-dimensional. Therefore the operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}$ is diagonal on odd lowest weight vectors Ψ_n^+ and Ψ_n^- but acts non-trivially on the two-dimensional subspace of even lowest weight vectors spanned on Φ_n^+ and Φ_n^-.
In matrix form we have:

$$\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)\begin{pmatrix} \Phi_n^+ \\ \Phi_n^- \\ \Psi_n^+ \\ \Psi_n^- \end{pmatrix} = \begin{pmatrix} A_n(u) & B_n(u) & 0 & 0 \\ C_n(u) & D_n(u) & 0 & 0 \\ 0 & 0 & F_n(u) & 0 \\ 0 & 0 & 0 & E_n(u) \end{pmatrix}\begin{pmatrix} \Phi_n^+ \\ \Phi_n^- \\ \Psi_n^+ \\ \Psi_n^- \end{pmatrix} \tag{4.3.1}$$

The matrix relation (2.2.5) leads to a set of recurrence relations for the coefficients $A_n, ..., E_n$ with the general solution [9]:

$$A_n(u) = (-1)^{n+1}\frac{\Gamma\left(\mathrm{u} + \ell_n\right)}{\Gamma\left(-\mathrm{u} + \ell_n + 1\right)} \cdot \tag{4.3.2}$$

$$\cdot\frac{(\ell_1 - b_1)(\ell_1 + b_1)\left(\mathrm{u} - b_1 - b_2\right) - \left(\mathrm{u} + b_1 + b_2\right)\left(\mathrm{u} - b_1 - \ell_2\right)\left(\mathrm{u} - b_1 + \ell_2\right)}{(\ell_2 - b_2)(\ell_1 + b_1)}$$

$$B_n(u) = (-1)^n\frac{\Gamma\left(\mathrm{u} + \ell_n\right)}{\Gamma\left(-\mathrm{u} + \ell_n\right)} \ ; \ C_n(u) = (-1)^n\frac{\Gamma\left(\mathrm{u} + \ell_n + 1\right)}{\Gamma\left(-\mathrm{u} + \ell_n + 1\right)} \cdot \frac{(\ell_2 + b_2)(\ell_1 - b_1)}{(\ell_2 - b_2)(\ell_1 + b_1)}$$

$$D_n(u) = (-1)^{n+1}\frac{\Gamma\left(\mathrm{u} + \ell_n + 1\right)}{\Gamma\left(-\mathrm{u} + \ell_n\right)} \cdot \frac{\mathrm{u} + b_2 - b_1}{(\ell_2 - b_2)(\ell_1 + b_1)}$$

$$E_n(u) = (-1)^n\frac{\Gamma\left(\mathrm{u} + \ell_n + 1\right)}{\Gamma\left(-\mathrm{u} + \ell_n + 1\right)} \cdot \frac{(\mathrm{u} + b_2 - \ell_1)(\mathrm{u} + b_2 + \ell_1)}{(\ell_2 - b_2)(\ell_1 + b_1)}$$

$$F_n(u) = (-1)^n \frac{\Gamma(\mathrm{u}+\ell_n+1)}{\Gamma(-\mathrm{u}+\ell_n+1)} \cdot \frac{(\mathrm{u}-b_1-\ell_2)(\mathrm{u}-b_1+\ell_2)}{(\ell_2-b_2)(\ell_1+b_1)}$$

where we used the notations:

$$\ell_n \equiv n+\ell_1+\ell_2 \ ; \ \mathrm{u} \equiv u+b_2-b_1.$$

As usual the obtained general solution of the Yang-Baxter equation is fixed up to overall normalization. We choose the normalization such that the R-matrix coincides with the permutation operator for $u=0$ and $\vec{\ell}_1=\vec{\ell}_2$. Note that the transformation from our basis vectors $\Phi_n^{\pm}$ to the ones diagonalizing the R-matrix depends on n and this does not result in simpler formulae. [1]

5 The operator $\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)$ as integral operator

Let us go to the derivation of the explicit representation for the $\mathbb{R}$-matrix as integral operator acting on polynomials $\psi(Z_1,Z_2) \in V_{\vec{\ell}_1} \otimes V_{\vec{\ell}_2}$

$$\left[\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)\psi\right](Z_1,Z_2) = \int\int \mathrm{d}Z_3\mathrm{d}Z_4\mathcal{R}(Z_1,Z_2|Z_3,Z_4)\psi(Z_3,Z_4) \tag{5.0.1}$$

where $Z \equiv \{z,\theta,\bar{\theta}\}$ and we use the standard superintegration

$$\int \mathrm{d}Z \equiv \int \mathrm{d}z \int \mathrm{d}\theta \int \mathrm{d}\bar{\theta}.$$

We suppose that it is possible to integrate by parts so that the equations (4.2.3), (4.2.4) give the set of differential equations for the kernel $\mathcal{R}(\vec{Z}) = \mathcal{R}(Z_1,Z_2|Z_3,Z_4)$

$$(\mathbb{R}_1+\mathbb{R}_2+\mathbb{R}_3+\mathbb{R}_4)R(\vec{Z}) = 0 \tag{5.0.2}$$

$$\left(\frac{u}{2}(\mathbb{R}_1-\mathbb{R}_2)+\mathbb{R}_1\mathbb{R}_2\right)R(\vec{Z}) = \left(\frac{u}{2}(\mathbb{R}_4-\mathbb{R}_3)+\mathbb{R}_4\mathbb{R}_3\right)R(\vec{Z}) \tag{5.0.3}$$

where the matrices $\mathbb{R}_i$, $i=3,4$ arise after integration by parts

$$\mathbb{R}_i(u) = \begin{pmatrix} S_i+B_i & -W_i^- & S_i^- \\ V_i^+ & 2B_i & V_i^- \\ S_i^+ & W_i^+ & B_i-S_i \end{pmatrix} \ ; \ \vec{\ell}_3 \equiv -\vec{\ell}_1 = (-\ell_1,-b_1) \ , \ \vec{\ell}_4 \equiv -\vec{\ell}_2 = (-\ell_2,-b_2)$$

---

[1] Note that we have admitted some inconsistency in notations in our previous work [9]. We have started from the general Yang-Baxter equation

$$\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u-v)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_3}(u)\mathbb{R}_{\vec{\ell}_2\vec{\ell}_3}(v) = \mathbb{R}_{\vec{\ell}_2\vec{\ell}_3}(v)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_3}(u)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u-v)$$

have choosen $\vec{\ell}_3=\vec{f}$ and obtained

$$\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u-v)\mathbb{R}_{\vec{\ell}_1\vec{f}}(u)\mathbb{R}_{\vec{\ell}_2\vec{f}}(v) = \mathbb{R}_{\vec{\ell}_2\vec{f}}(v)\mathbb{R}_{\vec{\ell}_1\vec{f}}(u)\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u-v)$$

Then we have "automatically" used the ansatz for $\mathbb{R}_{\vec{f}\vec{\ell}}$ instead of $\mathbb{R}_{\vec{\ell}\vec{f}}$. As a consequence the results for the matrix elements formulated there are referring actually to the operator $\mathbb{R}_{\vec{\ell}_2\vec{\ell}_1}(u)$

$$\mathbb{R}_{\vec{\ell}_2\vec{\ell}_1}(u) \equiv \mathbb{P}\mathbb{R}_{\vec{\ell}_1\vec{\ell}_2}(u)\mathbb{P}\Big|_{\vec{\ell}_1\leftrightarrow\vec{\ell}_2} \ ; \ \mathbb{P}\psi(Z_1,Z_2) \equiv \psi(Z_2,Z_1).$$

In the simplest $s\ell(2)$-case these matrices coincide $\mathbb{R}_{\ell_1\ell_2}(u) = \mathbb{R}_{\ell_2\ell_1}(u)$. In the case of superalgebra $s\ell(2|1)$ the situation is unusual because the flip-operator $\mathbb{P}$ acts nontrivially in the lowest weights basis we have used. The correct answer in selfconsistent notations is given in (4.3.2).

First we shall solve this system of differential equations and then specify selfconsistently the contours of integration.

Proposition 2 *The system of differential equations (5.0.2), (5.0.3) of second order is equivalent to the system of differential equations of first order*

$$\mathcal{D}_i\mathcal{R} = 0 \ , \ \mathcal{D}_i^{\pm}\mathcal{R} = 0 \ ; \ i = 1,2,3,4. \tag{5.0.4}$$

where the operators $\mathcal{D}_i, \mathcal{D}_i^{\pm}$ are defined in a following way

$$\begin{aligned}
\mathcal{D}_1 &\equiv \partial_1 - \frac{\bar\theta_{12}}{Z_{12}}D_1^- - \frac{\theta_{14}}{Z_{14}}D_1^+ + \frac{\mathrm{u}+\ell_1-\ell_2}{Z_{14}^+} - \frac{\mathrm{u}-\ell_1-\ell_2}{Z_{12}^-} + (\mathrm{u}-\ell_2-b_1)\frac{\theta_{14}\bar\theta_{12}}{Z_{14}Z_{12}} \\
\mathcal{D}_1^- &\equiv D_1^- - (\mathrm{u}-\ell_1-\ell_2)\frac{\theta_{12}}{Z_{12}} + (\mathrm{u}-\ell_2-b_1)\frac{\theta_{14}}{Z_{14}} \qquad (5.0.5) \\
\mathcal{D}_1^+ &\equiv D_1^+ + (\ell_1+b_1)\frac{\bar\theta_{14}}{Z_{14}} + (\mathrm{u}+b_2-b_1)\frac{Z_{43}^-}{Z_{41}^-}\left(\frac{\bar\theta_{43}}{Z_{43}} - \frac{\bar\theta_{23}}{Z_{23}}\right) \\
\mathcal{D}_2 &\equiv \partial_2 - \frac{\bar\theta_{23}}{Z_{23}}D_2^- - \frac{\theta_{21}}{Z_{21}}D_2^+ - \frac{\mathrm{u}-\ell_1-\ell_2}{Z_{21}^+} + \frac{\mathrm{u}-\ell_1+\ell_2}{Z_{23}^-} - (\mathrm{u}-\ell_1+b_2)\frac{\theta_{21}\bar\theta_{23}}{Z_{21}Z_{23}} \\
\mathcal{D}_2^+ &\equiv D_2^+ - (\mathrm{u}-\ell_1-\ell_2)\frac{\bar\theta_{21}}{Z_{21}} + (\mathrm{u}-\ell_1+b_2)\frac{\bar\theta_{23}}{Z_{23}} \qquad (5.0.6) \\
\mathcal{D}_2^- &\equiv D_2^- + (\ell_2-b_2)\frac{\theta_{23}}{Z_{23}} + (\mathrm{u}+b_2-b_1)\frac{Z_{34}^+}{Z_{32}^+}\left(\frac{\theta_{43}}{Z_{43}} - \frac{\theta_{14}}{Z_{14}}\right) \\
\mathcal{D}_3 &\equiv \partial_3 - \frac{\bar\theta_{32}}{Z_{32}}D_3^- - \frac{\theta_{34}}{Z_{34}}D_3^+ - \frac{\mathrm{u}+\ell_1+\ell_2}{Z_{34}^+} + \frac{\mathrm{u}-\ell_1+\ell_2}{Z_{32}^-} - (\mathrm{u}+\ell_2-b_1)\frac{\theta_{34}\bar\theta_{32}}{Z_{34}Z_{32}} \\
\mathcal{D}_3^+ &\equiv D_3^+ - (\mathrm{u}+\ell_1+\ell_2)\frac{\bar\theta_{34}}{Z_{34}} + (\mathrm{u}+\ell_2-b_1)\frac{\bar\theta_{32}}{Z_{32}} \qquad (5.0.7) \\
\mathcal{D}_3^- &\equiv D_3^- - (\ell_1-b_1)\frac{\theta_{23}}{Z_{23}} + (\mathrm{u}+b_2-b_1)\frac{Z_{21}^+}{Z_{23}^+}\left(\frac{\theta_{12}}{Z_{12}} - \frac{\theta_{14}}{Z_{14}}\right) \\
\mathcal{D}_4 &\equiv \partial_4 - \frac{\bar\theta_{43}}{Z_{43}}D_4^- - \frac{\theta_{41}}{Z_{41}}D_4^+ + \frac{\mathrm{u}+\ell_1-\ell_2}{Z_{41}^+} - \frac{\mathrm{u}+\ell_1+\ell_2}{Z_{43}^-} + (\mathrm{u}+\ell_1+b_2)\frac{\theta_{41}\bar\theta_{43}}{Z_{41}Z_{43}} \\
\mathcal{D}_4^- &\equiv D_4^- - (\mathrm{u}+\ell_1+\ell_2)\frac{\theta_{43}}{Z_{43}} + (\mathrm{u}+\ell_1+b_2)\frac{\theta_{41}}{Z_{41}} \qquad (5.0.8) \\
\mathcal{D}_4^+ &\equiv D_4^+ - (\ell_2+b_2)\frac{\bar\theta_{41}}{Z_{41}} + (\mathrm{u}+b_2-b_1)\frac{Z_{12}^-}{Z_{14}^-}\left(\frac{\bar\theta_{12}}{Z_{12}} - \frac{\bar\theta_{23}}{Z_{23}}\right)
\end{aligned}$$

where

$$D_i^+ = -\partial_{\theta_i} + \frac{1}{2}\bar\theta_i\partial_i \ ; \ D_i^- = -\partial_{\bar\theta_i} + \frac{1}{2}\theta_i\partial_i$$

$$Z_{ik}^{\pm} = Z_{ik} \pm \frac{1}{2}\theta_{ik}\bar\theta_{ik} \ ; \ D_i^{\pm}Z_{ik}^{\pm} = 0 \ ; \ D_k^{\mp}Z_{ik}^{\pm} = 0.$$

This system of twelve equations is invariant with respect the following discrete symmetry transformations:

$$1 \leftrightarrow 4 \ , \ 2 \leftrightarrow 3 \ ; \ (\ell_1, b_1) \leftrightarrow (-\ell_2, -b_2),$$

$$1 \leftrightarrow 2 \ , \ 3 \leftrightarrow 4 \ ; \ \theta \leftrightarrow \bar\theta \ , \ D^+ \leftrightarrow D^- \ ; \ (\ell_1, b_1) \leftrightarrow (\ell_2, -b_2).$$

5.1 Differential equations

We use the same trick as in the $s\ell(2)$-case and derive the simple first order differential equations for the kernel of R-operator. The matrix $\mathbb{R}$ has a very special structure

$$\mathbb{R} = \begin{pmatrix} 1 \\ -\bar\theta \\ z+\frac{1}{2}\theta\bar\theta \end{pmatrix} \begin{pmatrix} z-\frac{1}{2}\theta\bar\theta & -\theta & -1 \end{pmatrix} \partial - \begin{pmatrix} 1 \\ -\bar\theta \\ z+\frac{1}{2}\theta\bar\theta \end{pmatrix} \begin{pmatrix} \bar\theta & -1 & 0 \end{pmatrix} D^- -$$

$$- \begin{pmatrix} 0 \\ -1 \\ \theta \end{pmatrix} \begin{pmatrix} z-\frac{1}{2}\theta\bar\theta & \theta & -1 \end{pmatrix} D^+ + (\ell+b) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ z-\frac{1}{2}\theta\bar\theta & -\theta & 0 \end{pmatrix} + (\ell-b) \begin{pmatrix} 0 & 0 & 0 \\ -\bar\theta & -1 & 0 \\ z+\frac{1}{2}\theta\bar\theta & 0 & -1 \end{pmatrix}$$

where

$$D^+ = -\partial_\theta + \frac{1}{2}\bar\theta\partial \ ; \ D^- = -\partial_{\bar\theta} + \frac{1}{2}\theta\partial,$$

so that the matrix $\mathbb{R}f(Z)$, where $f(Z)$ is an even function, has two evident eigenvectors

$$\begin{pmatrix} z-\frac{1}{2}\theta\bar\theta & -\theta & -1 \end{pmatrix} \mathbb{R}f = -(\ell-b)f \cdot \begin{pmatrix} z-\frac{1}{2}\theta\bar\theta & -\theta & -1 \end{pmatrix}$$

$$\mathbb{R}f \begin{pmatrix} 1 \\ -\bar\theta \\ z+\frac{1}{2}\theta\bar\theta \end{pmatrix} = (\ell+b)f \cdot \begin{pmatrix} 1 \\ -\bar\theta \\ z+\frac{1}{2}\theta\bar\theta \end{pmatrix}$$

and all derivatives $\partial f, D^+f, D^-f$ disappear from the relations. Moreover, there are two other vectors with the properties

$$\begin{pmatrix} \theta & 1 & 0 \end{pmatrix} \mathbb{R}f = 2bf \cdot \begin{pmatrix} \theta & 1 & 0 \end{pmatrix} + \begin{pmatrix} z-\frac{1}{2}\theta\bar\theta & \theta & -1 \end{pmatrix} D^+f$$

$$\mathbb{R}f \begin{pmatrix} 0 \\ 1 \\ -\theta \end{pmatrix} = 2bf \cdot \begin{pmatrix} 0 \\ 1 \\ -\theta \end{pmatrix} + \begin{pmatrix} 1 \\ -\bar\theta \\ z+\frac{1}{2}\theta\bar\theta \end{pmatrix} D^-f.$$

We shall derive the first order differential equations for the kernel $\mathcal{R}(\vec{Z})$ by multiplying the equations (4.2.3), (4.2.4) by the appropriate vectors from the left and from the right. Let us start from the three variants for the choose of the such vectors. First we obtain the system of two first-order differential equations for the function $\mathcal{R}(\vec{Z})$

$$\mathbb{D}_1\mathcal{R} + \mathbb{D}_3\mathcal{R} = (\ell_2 - b_2 - \ell_4 + b_4)Z^+_{24}\mathcal{R}$$

$$\left(\frac{u}{2} + \ell_2 + b_2\right)\mathbb{D}_1\mathcal{R} + \left(\frac{u}{2} + \ell_4 - b_4\right)\mathbb{D}_3\mathcal{R} = -\frac{u}{2}(\ell_2 - \ell_4 + b_2 + b_4)Z^+_{24}\mathcal{R}$$

which is equivalent to the system

$$\mathbb{D}_1\mathcal{R} = -\left(u + \ell_4 - b_4\right)Z^+_{24}\mathcal{R} \ ; \ \mathbb{D}_3\mathcal{R} = \left(u + \ell_2 + b_2\right)Z^+_{24}\mathcal{R}. \tag{5.1.9}$$

We use the notation

$$\mathbb{D}_k \equiv \begin{pmatrix} z_4-\frac{1}{2}\theta_4\bar\theta_4 & -\theta_4 & -1 \end{pmatrix} \mathbb{R}_k \begin{pmatrix} 1 \\ -\bar\theta_2 \\ z_2+\frac{1}{2}\theta_2\bar\theta_2 \end{pmatrix} =$$

$$= -Z^-_{k2}Z^+_{k4}\partial_k + Z^+_{k4}\bar\theta_{k2}D^-_k + Z^-_{k2}\theta_{k4}D^+_k - (\ell_k + b_k)Z^+_{k4} - (\ell_k - b_k)Z^-_{k2} + 2b_k\theta_{k4}\bar\theta_{k2} \ ; \ k = 1,3$$

for the arising differential operators. Secondly we obtain the system of differential equations

$$\mathbb{B}_1\mathcal{R} + \mathbb{B}_3\mathcal{R} = (\ell_4 - b_4 - 2b_2)\theta_{24}\mathcal{R} + Z^+_{24}D^-_2\mathcal{R}$$

$$\left(\frac{u}{2} + 2b_2\right)\mathbb{B}_1\mathcal{R} + \left(\frac{u}{2} + \ell_4 - b_4\right)\mathbb{B}_3\mathcal{R} = -\frac{u}{2}(\ell_4 - b_4 - 2b_2)\theta_{24}\mathcal{R} - \frac{u}{2}Z^+_{24}D^-_2\mathcal{R} - \mathbb{D}_1 D^-_2\mathcal{R}$$

which is equivalent to the system

$$\mathbb{B}_1\mathcal{R} = (u + \ell_4 - b_4)\,\theta_{24}\mathcal{R}\ ;\ \mathbb{B}_3\mathcal{R} = -\,(u + 2b_2)\,\theta_{24}\mathcal{R} + Z^+_{24}D^-_2\mathcal{R}, \tag{5.1.10}$$

by conditions (5.1.9) and where

$$\mathbb{B}_k \equiv \begin{pmatrix} z_4 - \frac{1}{2}\theta_4\bar{\theta}_4 & -\theta_4 & -1 \end{pmatrix}\mathbb{R}_k \begin{pmatrix} 0 \\ 1 \\ -\theta_2 \end{pmatrix} =$$

$$= Z^+_{k4}\theta_{k2}\partial_k - Z^+_{k4}D^-_k - \theta_{k2}\theta_{k4}D^+_k + (\ell_k + b_k)\theta_{k2} - (\ell_k - b_k)\theta_{24}\ ;\ k = 1, 3.$$

The last system is analogous to the previous ones:

$$\mathbb{C}_1\mathcal{R} + \mathbb{C}_3\mathcal{R} = (\ell_2 + b_2 + 2b_4)\bar{\theta}_{24}\mathcal{R} + Z^+_{24}D^+_4\mathcal{R}$$

$$\left(\frac{u}{2} + \ell_2 + b_2\right)\mathbb{C}_1\mathcal{R} + \left(\frac{u}{2} + 1 - 2b_4\right)\mathbb{C}_3\mathcal{R} = -\frac{u}{2}(\ell_2 + b_2 + 2b_4)\bar{\theta}_{24}\mathcal{R} - \frac{u}{2}Z^+_{24}D^+_4\mathcal{R} + D^+_4\mathbb{D}_3\mathcal{R}$$

and is equivalent to the system

$$\mathbb{C}_3\mathcal{R} = (u + \ell_2 + b_2)\,\bar{\theta}_{24}\mathcal{R}\ ;\ \mathbb{C}_1\mathcal{R} = -\,(u - 2b_4)\,\bar{\theta}_{24}\mathcal{R} + Z^+_{24}D^+_4\mathcal{R}, \tag{5.1.11}$$

by conditions (5.1.9). The operators $\mathbb{C}_k$ are

$$\mathbb{C}_k \equiv \begin{pmatrix} \bar{\theta}_4 & 1 & 0 \end{pmatrix}\mathbb{R}_k \begin{pmatrix} 1 \\ -\bar{\theta}_2 \\ z_2 + \frac{1}{2}\theta_2\bar{\theta}_2 \end{pmatrix} =$$

$$= -Z^-_{k2}\bar{\theta}_{k4}\partial_k + \bar{\theta}_{k4}\bar{\theta}_{k2}D^-_k + Z^-_{k2}D^+_k - (\ell_k + b_k)\bar{\theta}_{24} - (\ell_k - b_k)\bar{\theta}_{k2}\ ;\ k = 1, 3.$$

The obtained system of the six equations can be transformed to the equivalent form. The $\mathbb{D}_{1,3}$-equations (5.1.9) are equivalent to the equations $\mathcal{D}_1\mathcal{R} = 0$, $\mathcal{D}_3\mathcal{R} = 0$. The $\mathbb{B}_1$-equation from (5.1.10) can be transformed to the equation $\mathcal{D}^-_1\mathcal{R} = 0$ by the condition $\mathcal{D}_1\mathcal{R} = 0$. The $\mathbb{C}_3$-equation from (5.1.11) can be transformed to the equation $\mathcal{D}^+_3\mathcal{R} = 0$ by the condition $\mathcal{D}_3\mathcal{R} = 0$. The remaining $\mathbb{B}_3, \mathbb{C}_1$-equations are equivalent to the more involved equations

$$-Z^+_{34}\left(1 - \frac{\theta_{32}\bar{\theta}_{32}}{Z_{32}}\right)\left(D^-_3 + (\mathrm{u} + \ell_2 - \ell_1)\frac{\theta_{32}}{Z_{32}} - (\mathrm{u} + \ell_2 - b_1)\frac{\theta_{34}}{Z_{34}}\right)\mathcal{R} = Z^+_{24}D^-_2\mathcal{R} + (\ell_2 - b_2)\theta_{24}\mathcal{R}$$

$$Z^-_{12}\left(1 + \frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)\left(D^+_1 + (\mathrm{u} + \ell_1 - \ell_2)\frac{\bar{\theta}_{14}}{Z_{14}} - (\mathrm{u} - \ell_2 - b_1)\frac{\bar{\theta}_{12}}{Z_{12}}\right)\mathcal{R} = Z^+_{24}D^+_4\mathcal{R} - (\ell_2 + b_2)\bar{\theta}_{24}\mathcal{R}$$

by the conditions $\mathcal{D}_1\mathcal{R} = 0$, $\mathcal{D}_3\mathcal{R} = 0$.

It is possible to derive the mirror system of six equations in the same manner by choosing appropriate vectors for multiplication from the left and from the right. In fact, the resulting equations can be obtained from the present ones by the symmetry transformation

$$1 \to 4\ ,\ 2 \to 3\ ;\ (\ell_1, b_1) \to (\ell_4, b_4) = (-\ell_2, -b_2)\ ,\ (\ell_2, b_2) \to (\ell_3, b_3) = (-\ell_1, -b_1)$$

In this way we obtain the equations $\mathcal{D}_4\mathcal{R} = 0$, $\mathcal{D}_2\mathcal{R} = 0$ and $\mathcal{D}_4^-\mathcal{R} = 0$, $\mathcal{D}_3^+\mathcal{R} = 0$ and the mirror partners of the more involved equations:

$$-Z_{21}^+\left(1+\frac{\theta_{32}\bar{\theta}_{32}}{Z_{32}}\right)\left(D_2^- + (\mathrm{u}+\ell_2-\ell_1)\frac{\theta_{32}}{Z_{32}} - (\mathrm{u}-\ell_1+b_2)\frac{\theta_{21}}{Z_{21}}\right)\mathcal{R} = Z_{31}^+ D_3^-\mathcal{R} - (\ell_1-b_1)\theta_{31}\mathcal{R}$$

$$Z_{43}^-\left(1-\frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)\left(D_4^+ + (\mathrm{u}+\ell_1-\ell_2)\frac{\bar{\theta}_{14}}{Z_{14}} - (\mathrm{u}+\ell_1+b_2)\frac{\bar{\theta}_{43}}{Z_{43}}\right)\mathcal{R} = Z_{31}^+ D_1^+\mathcal{R} + (\ell_1+b_1)\bar{\theta}_{31}\mathcal{R}.$$

The system of four involved equations decomposes into two independent systems of two equations. It can be shown that the system which involves the covariant derivatives D_1^+ and D_4^+ is equivalent to the system of the equations $\mathcal{D}_1^+\mathcal{R} = 0$, $\mathcal{D}_4^+\mathcal{R} = 0$ and the system which involves the covariant derivatives D_2^- and D_3^- is equivalent to the system of the equations $\mathcal{D}_2^-\mathcal{R} = 0$, $\mathcal{D}_3^-\mathcal{R} = 0$.

The next step is to prove the reverse, that the system of equations (5.0.2), (5.0.3) is the consequence of the system (5.0.4). It is easy to rewrite the $s\ell(2|1)$-generators and therefore the equations of $s\ell(2)$-invariance (5.0.2) in terms of introduced operators $\mathcal{D}_i, \mathcal{D}_i^{\pm}$:

$$\mathbb{S}^- = -\sum_{i=1}^4 \mathcal{D}_i \ ; \ \mathbb{V}^- = \sum_{i=1}^4 \bar{\theta}_i\mathcal{D}_i - \mathcal{D}_i^+ \ ; \ \mathbb{W}^- = \sum_{i=1}^4 \theta_i\mathcal{D}_i - \mathcal{D}_i^-$$

$$\mathbb{S}^+ = \sum_{i=1}^4 z_i^2\mathcal{D}_i - z_i\theta_i\mathcal{D}_i^+ - z_i\bar{\theta}_i\mathcal{D}_i^- \ ; \ \mathbb{V}^+ = \sum_{i=1}^4 -z_i\bar{\theta}_i\mathcal{D}_i + \left(z_i - \frac{1}{2}\theta_i\bar{\theta}_i\right)\mathcal{D}_i^+$$

$$\mathbb{W}^+ = \sum_{i=1}^4 -z_i\theta_i\mathcal{D}_i + \left(z_i + \frac{1}{2}\theta_i\bar{\theta}_i\right)\mathcal{D}_i^-$$

$$\mathbb{S} = \sum_{i=1}^4 z_i\mathcal{D}_i - \frac{1}{2}\bar{\theta}_i\mathcal{D}_i^- - \frac{1}{2}\theta_i\mathcal{D}_i^+ \ ; \ \mathbb{B} = -\frac{1}{2}\sum_{i=1}^4 \theta_i\bar{\theta}_i\mathcal{D}_i + \bar{\theta}_i\mathcal{D}_i^- - \theta_i\mathcal{D}_i^+$$

The last step is absolutely analogous to the $s\ell(2)$-case. Due to $s\ell(2|1)$-invariance we can use one K_{13}-equation in (5.0.3) and rewrite it in terms of operators $\mathcal{D}_i, \mathcal{D}_i^{\pm}$. We omit these formulae for simplicity.

5.2 Solution of the system of differential equations

It is useful to transform the main system of equations (5.0.5)-(5.0.8) to a simpler homogeneous form. We shall use the following ansatz for this purpose

$$\mathcal{R}(\vec{Z}) = Z_{12}^a Z_{14}^b Z_{32}^c Z_{34}^d \cdot \left(1+\frac{\theta_{12}\bar{\theta}_{12}}{Z_{12}}\right)^{\alpha}\left(1+\frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)^{\beta}\left(1+\frac{\theta_{32}\bar{\theta}_{32}}{Z_{32}}\right)^{\gamma}\left(1+\frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)^{\delta} R(\vec{Z})$$

$$a \equiv \mathrm{u}-\ell_1-\ell_2 \ ; \ b \equiv -\mathrm{u}-\ell_1+\ell_2 \ ; \ c \equiv -\mathrm{u}+\ell_1-\ell_2 \ ; \ d \equiv \mathrm{u}+\ell_1+\ell_2$$

$$\beta = b_1-\alpha \ ; \ \gamma = -b_2-\alpha \ ; \ \delta = b_2-b_1+\alpha.$$

Proposition 3 *The system of differential equations (5.0.5)-(5.0.8) for the function $\mathcal{R}(\vec{Z})$ is transformed to the simpler system of differential equations for the function $R(\vec{Z})$*

$$\mathcal{D}_i R(\vec{Z}) = 0 \ , \ \mathcal{D}_i^{\pm} R(\vec{Z}) = 0 \ ; \ i = 1,2,3,4. \tag{5.2.12}$$

where operators $\mathcal{D}_i, \mathcal{D}_i^{\pm}$ are defined in a following way

$$\begin{aligned}
\mathcal{D}_1 &\equiv \partial_1 - \frac{\bar{\theta}_{12}}{Z_{12}} D_1^- - \frac{\theta_{14}}{Z_{14}} D_1^+ \; ; \; \mathcal{D}_1^- \equiv D_1^- - A\left(\frac{\theta_{12}}{Z_{12}} - \frac{\theta_{14}}{Z_{14}}\right) \\
\mathcal{D}_1^+ &\equiv D_1^+ + A\left(\frac{\bar{\theta}_{12}}{Z_{12}} - \frac{\bar{\theta}_{14}}{Z_{14}}\right) + (A+B)\frac{Z_{43}^-}{Z_{41}^-}\left(\frac{\bar{\theta}_{43}}{Z_{43}} - \frac{\bar{\theta}_{23}}{Z_{23}}\right) \\
\mathcal{D}_2 &\equiv \partial_2 - \frac{\bar{\theta}_{23}}{Z_{23}} D_2^- - \frac{\theta_{21}}{Z_{21}} D_2^+ \; ; \; \mathcal{D}_2^+ \equiv D_2^+ - A\left(\frac{\bar{\theta}_{21}}{Z_{21}} - \frac{\bar{\theta}_{23}}{Z_{23}}\right) \\
\mathcal{D}_2^- &\equiv D_2^- + A\left(\frac{\theta_{21}}{Z_{21}} - \frac{\theta_{23}}{Z_{23}}\right) + (A+B)\frac{Z_{34}^+}{Z_{32}^+}\left(\frac{\theta_{43}}{Z_{43}} - \frac{\theta_{14}}{Z_{14}}\right) \\
\mathcal{D}_3 &\equiv \partial_3 - \frac{\bar{\theta}_{32}}{Z_{32}} D_3^- - \frac{\theta_{34}}{Z_{34}} D_3^+ \; ; \; \mathcal{D}_3^+ \equiv D_3^+ - B\left(\frac{\bar{\theta}_{34}}{Z_{34}} - \frac{\bar{\theta}_{32}}{Z_{32}}\right) \\
\mathcal{D}_3^- &\equiv D_3^- + B\left(\frac{\theta_{34}}{Z_{34}} - \frac{\theta_{32}}{Z_{32}}\right) + (A+B)\frac{Z_{21}^+}{Z_{23}^+}\left(\frac{\theta_{12}}{Z_{12}} - \frac{\theta_{14}}{Z_{14}}\right) \\
\mathcal{D}_4 &\equiv \partial_4 - \frac{\bar{\theta}_{43}}{Z_{43}} D_4^- - \frac{\theta_{41}}{Z_{41}} D_4^+ \; ; \; \mathcal{D}_4^- \equiv D_4^- - B\left(\frac{\theta_{43}}{Z_{43}} - \frac{\theta_{41}}{Z_{41}}\right) \\
\mathcal{D}_4^+ &\equiv D_4^+ + B\left(\frac{\bar{\theta}_{43}}{Z_{43}} - \frac{\bar{\theta}_{41}}{Z_{41}}\right) + (A+B)\frac{Z_{12}^-}{Z_{14}^-}\left(\frac{\bar{\theta}_{12}}{Z_{12}} - \frac{\bar{\theta}_{23}}{Z_{23}}\right)
\end{aligned} \tag{5.2.13}$$

where

$$A \equiv \frac{\mathrm{u} - \ell_1 - \ell_2}{2} - \alpha \; ; \; B \equiv \frac{\mathrm{u} + \ell_1 + \ell_2}{2} + b_2 - b_1 + \alpha \; ; \; A + B = \mathrm{u} + b_2 - b_1$$

The common solution of these twelve equations $\mathcal{D}_i\mathcal{R} = 0, \mathcal{D}_i^{\pm}\mathcal{R} = 0 \;\; i = 1,2,3,4$ is the manifestly superconformal-invariant function (see Appendix A):

$$\begin{aligned}
R(\vec{Z}) &= \left(1 + A\frac{\theta_{12}^3\bar{\theta}_{12}^3}{Z_{12}^3}\right)\cdot\left(1 + B\frac{\theta_{14}^3\bar{\theta}_{14}^3}{Z_{14}^3}\right) - (A+B)\frac{\theta_{14}^3\bar{\theta}_{12}^3}{Z_{14}^3} = \\
&= \left(1 + \frac{\theta_{12}^3\bar{\theta}_{12}^3}{Z_{12}^3} - \frac{\theta_{14}^3\bar{\theta}_{12}^3}{Z_{14}^3}\right)^A \cdot \left(1 + \frac{\theta_{14}^3\bar{\theta}_{14}^3}{Z_{14}^3} - \frac{\theta_{14}^3\bar{\theta}_{12}^3}{Z_{14}^3}\right)^B
\end{aligned} \tag{5.2.14}$$

where

$$Z_{ij}^k \equiv \frac{Z_{ij}}{Z_{ik}Z_{jk}} \; , \; \theta_{ij}^k \equiv \frac{\theta_{ik}}{Z_{ik}} - \frac{\theta_{jk}}{Z_{jk}} \; , \; \bar{\theta}_{ij}^k \equiv \frac{\bar{\theta}_{ik}}{Z_{ik}} - \frac{\bar{\theta}_{jk}}{Z_{jk}}$$

This proposition can be proved by direct calculations. It is remarkable that all involved operators $\mathcal{D}_i$, $\mathcal{D}_i^{\pm}$ can be obtained by the similarity transformation from the simplest derivatives $\partial_i, D_i^{\pm}$

$$\mathcal{D}_i = R\partial_i R^{-1} \; ; \; \mathcal{D}_i^{\pm} = R D_i^{\pm} R^{-1} \; ; \; R = R(\vec{Z}).$$

5.3 The integral representation.

It remains to fix the contours of integrations. The $\mathbb{R}$-operator acts on the space of polynomials $\psi(Z_1, Z_2)$ which can be of arbitrary large degree. The requirement of convergence allows the compact contours only. The final representation for the R-matrix is:

$$\left[\mathbb{R}_{\vec{\ell}_1,\vec{\ell}_2}(u)\psi\right](Z_1, Z_2) = \rho \int_{z_1}^{z_2} \mathrm{d}Z_3 \int_{z_1}^{z_3} \mathrm{d}Z_4 \mathcal{R}(Z_1, Z_2|Z_3, Z_4)\psi(Z_3, Z_4) \tag{5.3.15}$$

where $Z \equiv \{z, \theta, \bar{\theta}\}$

$$\int_{z_1}^{z_2} \mathrm{d}Z_3 \equiv \int_{z_1}^{z_2} \mathrm{d}z_3 \int \mathrm{d}\theta_3 \int \mathrm{d}\bar{\theta}_3. \ ; \ \int_{z_1}^{z_3} \mathrm{d}Z_4 \equiv \int_{z_1}^{z_3} \mathrm{d}z_4 \int \mathrm{d}\theta_4 \int \mathrm{d}\bar{\theta}_4$$

and

$$\mathcal{R}(\vec{Z}) = Z_{12}^a Z_{14}^b Z_{32}^c Z_{34}^d \cdot \left(1 + \frac{\theta_{12}\bar{\theta}_{12}}{Z_{12}}\right)^\alpha \left(1 + \frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)^\beta \left(1 + \frac{\theta_{32}\bar{\theta}_{32}}{Z_{32}}\right)^\gamma \left(1 + \frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)^\delta \cdot$$

$$\cdot \left\{ \left(1 + A\frac{\theta_{12}^3\bar{\theta}_{12}^3}{Z_{12}^3}\right) \cdot \left(1 + B\frac{\theta_{14}^3\bar{\theta}_{14}^3}{Z_{14}^3}\right) - (A + B)\frac{\theta_{14}^3\bar{\theta}_{12}^3}{Z_{14}^3} \right\} \tag{5.3.16}$$

$$a \equiv \mathrm{u} - \ell_1 - \ell_2 \ ; \ b \equiv -\mathrm{u} - \ell_1 + \ell_2 \ ; \ c \equiv -\mathrm{u} + \ell_1 - \ell_2 \ ; \ d \equiv \mathrm{u} + \ell_1 + \ell_2$$

$$\beta = b_1 - \alpha \ ; \ \gamma = -b_2 - \alpha \ ; \ \delta = b_2 - b_1 + \alpha.$$

The normalisation factor ρ is chosen so that $\mathbb{R} : 1 \mapsto 1$

$$\rho = \frac{\Gamma(c + b + d)}{\Gamma(b)\Gamma(c)\Gamma(d)} \frac{(\mathrm{u} - \ell_1 + b_2)(\mathrm{u} - \ell_2 - b_1)(\mathrm{u} + \ell_1 + \ell_2)}{-\mathrm{u} + \ell_1 + \ell_2} \tag{5.3.17}$$

It is possible to check *a posteriori* that the obtained integral operator has the matrix elements (4.3.1). Using the formulae from Appendix B we derive the formulae for the even basis vectors:

$$\int_{z_1}^{z_2} \mathrm{d}Z_3 \int_{z_1}^{z_3} \mathrm{d}Z_4 \mathcal{R}(Z_1, Z_2|Z_3, Z_4) \left(Z_{34} + \frac{\theta_{34}\bar{\theta}_{34}}{2}\right)^n =$$

$$= \frac{\Gamma(b)\Gamma(c)\Gamma(d + n)}{\Gamma(c + b + d + n)} \cdot \left\{ \mathcal{R}_{++} \left(Z_{21} + \frac{\theta_{21}\bar{\theta}_{21}}{2}\right)^n + \mathcal{R}_{-+} \left(Z_{21} - \frac{\theta_{21}\bar{\theta}_{21}}{2}\right)^n \right\}$$

where

$$\mathcal{R}_{++} = (\ell_1 + b_1)(\ell_2 - b_2) \ ; \ \mathcal{R}_{-+} = \frac{\mathrm{u}^3 + \mathrm{u}^2(b_2 - b_1) - \mathrm{u}(\ell_1^2 + \ell_2^2 + 2b_1 b_2) + (b_1 + b_2)(\ell_1^2 - \ell_2^2)}{-\mathrm{u} + \ell_1 + \ell_2 + n}$$

and

$$\int_{z_1}^{z_2} \mathrm{d}Z_3 \int_{z_1}^{z_3} \mathrm{d}Z_4 \mathcal{R}(Z_1, Z_2|Z_3, Z_4) \left(Z_{34} - \frac{\theta_{34}\bar{\theta}_{34}}{2}\right)^n =$$

$$= \frac{\Gamma(b)\Gamma(c)\Gamma(d + n)}{\Gamma(c + b + d + n)} \cdot \left\{ \mathcal{R}_{+-} \left(Z_{21} + \frac{\theta_{21}\bar{\theta}_{21}}{2}\right)^n + \mathcal{R}_{--} \left(Z_{21} - \frac{\theta_{21}\bar{\theta}_{21}}{2}\right)^n \right\}$$

where

$$\mathcal{R}_{+-} = -(\mathrm{u} + b_2 - b_1)(\mathrm{u} + \ell_1 + \ell_2 + n) \ ; \ \mathcal{R}_{--} = \frac{(\ell_1 - b_1)(\ell_2 + b_2)(\mathrm{u} + \ell_1 + \ell_2 + n)}{-\mathrm{u} + \ell_1 + \ell_2 + n}$$

$$\mathcal{R}_{++} + \mathcal{R}_{-+}|_{n=0} = \mathcal{R}_{+-} + \mathcal{R}_{--}|_{n=0} = \frac{-\mathrm{u} + \ell_1 + \ell_2}{(\mathrm{u} - \ell_1 + b_2)(\mathrm{u} - \ell_2 - b_1)(\mathrm{u} + \ell_1 + \ell_2)}$$

The analogous formulae for the odd vectors are much simpler:

$$\int_{z_1}^{z_2} \mathrm{d}Z_3 \int_{z_1}^{z_3} \mathrm{d}Z_4 \mathcal{R}(Z_1, Z_2|Z_3, Z_4) Z_{34}^n \cdot \begin{pmatrix} \theta_{34} \\ \bar{\theta}_{34} \end{pmatrix} =$$

$$= -\frac{\Gamma(b)\Gamma(c)\Gamma(d+n+1)}{\Gamma(c+b+d+n+1)} Z_{21}^n \cdot \begin{pmatrix} (\mathrm{u} + b_2 + \ell_1)(\mathrm{u} + b_2 - \ell_1)\theta_{21} \\ (\mathrm{u} - b_1 - \ell_2)(\mathrm{u} - b_1 + \ell_2)\bar{\theta}_{21} \end{pmatrix}$$

These formulae for the action of the R-matrix on the basis vectors coincide up to overall normalisation with the ones (4.3.1) obtained in our previous paper [9].

6 Conclusions

We have derived the integral operator form of the rational $sl(2|1)$ symmetric solution of the Yang-Baxter equation acting on the tensor product of two general lowest weight representations $\vec{\ell_1} = (\ell_1, b_1)$ and $\vec{\ell_2} = (\ell_2, b_2)$. The lowest weight representations space is identified with the space of polynomials in one even (z) and two odd ($\theta, \bar{\theta}$). The kernel of the R-operator is a superconformal four-point function of primary fields with weights $\vec{\ell_1}, \vec{\ell_2}$ and $\vec{\ell_3} = -\vec{\ell_1}$, $\vec{\ell_4} = -\vec{\ell_2}$. This function of the supercoordinates of four points arises as a simple sum of terms in powers of $Z_{ij}, \theta_{ij}, \bar{\theta}_{ij}$. The simplicity of the result makes this integral operator form useful in applications and further investigations.

Following the well known procedure, the general R-operator is derived from the Yang-Baxter equation involving the latter and the known R-operator $R_{f,\ell}$ acting in the tensor product of the fundamental representation (in matrix form) and an arbitrary representation. We have formulated the conditions involved in this Yang-Baxter relation as differential equations on the kernel.

There is a remarkable reductions of these equations, being an overdetermined system of second order equations, to a simple set of first order equations. In this way the obtained kernel expresses the similarity transformation relating this consistent system of first order equations to the trivial one being the condition for a function to be constant in all variables $z_i, \theta_i, \bar{\theta}_i$. We have obtained this reduction by observing that the Lax matrix representing $R_{f,\ell}$ has a peculiar structure implying that appropriate projections reduce the number of derivatives.

We have calculated the matrix elements of the obtained R-operator. The result is consistent with the one obtained earlier from recurrence relations in the framework of the matrix formulation. We have quoted the formulae to be used for calculating the action of the integral operator.

The method of calculation relying on this reduction seems to be general enough to become useful for solving the Yang-Baxter equations with other symmetries. The structure on which the reduction relies deserves further study.

7 Acknowledgments

We thank G.Korchemsky and A.Manashov for the stimulating discussion and critical remarks. This work has been supported by Deutsche Forschungsgemeinschaft, grant No KI

623/1-2 and by INTAS,grant No 96-524. One of us (D.K.) is grateful to the German Ministry BMBF for support. The work of S.D. was partially supported by RFFI-grant No 00-01-005000.

Appendix A

In this Appendix we shall construct the superconformal invariant N-point function with zero global U(1)-charge. The requirement of $s\ell(2|1)$-invariance (5.0.2) results in the equations:

$$V^-\mathcal{R}(\vec{Z})=0\ ;\ W^-\mathcal{R}(\vec{Z})=0\ ;\ V^+\mathcal{R}(\vec{Z})=0\ ;\ W^+\mathcal{R}(\vec{Z})=0$$

where $V^-=\sum_{i=1}^{4}V_i^-$ and so on. Note we show the independent equations only. The others follow from the commutation relations

$$\{V^\pm,W^\pm\}=\pm S^\pm\ ;\ \{V^\pm,W^\mp\}=-S\pm B,$$

Let us consider the general case of N-point functions with zero total $U(1)$-charge: $\vec{Z}=(Z_1,...,Z_N)\ ;\ \sum_i b_i=0$. It is convenient to use the global $s\ell(2|1)$-transformations which are generated by the lowering operators V^-,W^-

$$e^{\epsilon V^-}\Phi(z;\theta,\bar\theta)=\Phi\left(z+\frac{\epsilon\bar\theta}{2};\theta+\epsilon,\bar\theta\right)\ ;\ e^{\epsilon W^-}\Phi(z;\theta,\bar\theta)=\Phi\left(z+\frac{\epsilon\theta}{2};\theta,\bar\theta+\epsilon\right).$$

and the rising operators V^+,W^+

$$e^{\epsilon V^+}\Phi(z;\theta,\bar\theta)=\frac{1}{(1+\epsilon\bar\theta)^{\ell-b}}\Phi\left(\frac{z}{1+\frac{\epsilon\bar\theta}{2}};\frac{\theta-\epsilon z}{1+\frac{\epsilon\bar\theta}{2}},\bar\theta\right),$$

$$e^{\epsilon W^+}\Phi(z;\theta,\bar\theta)=\frac{1}{(1+\epsilon\theta)^{\ell+b}}\Phi\left(\frac{z}{1+\frac{\epsilon\theta}{2}};\theta,\frac{\bar\theta-\epsilon z}{1+\frac{\epsilon\theta}{2}}\right),$$

so that the requirement of superconformal invariance results in the system of equations:

$$V^-:\ \ \mathcal{R}(Z_1...Z_N)=\mathcal{R}\left(z_1+\frac{\epsilon\bar\theta_1}{2},\theta_1+\epsilon,\bar\theta_1...z_N+\frac{\epsilon\bar\theta_N}{2},\theta_N+\epsilon,\bar\theta_N\right)$$

$$W^-:\ \ \mathcal{R}(Z_1...Z_N)=\mathcal{R}\left(z_1+\frac{\epsilon\theta_1}{2},\theta_1,\bar\theta_1+\epsilon...z_N+\frac{\epsilon\theta_N}{2},\theta_N,\bar\theta_N+\epsilon\right)$$

$$V^+:\ \ \mathcal{R}(Z_1...Z_N)=\prod_{i=1}^{N}\frac{1}{(1+\epsilon\bar\theta_i)^{\ell_i-b_i}}\mathcal{R}\left(\frac{z_1}{1+\frac{\epsilon\bar\theta_1}{2}},\frac{\theta_1-\epsilon z_1}{1+\frac{\epsilon\bar\theta_1}{2}},\bar\theta_1...\frac{z_N}{1+\frac{\epsilon\bar\theta_N}{2}},\frac{\theta_N-\epsilon z_N}{1+\frac{\epsilon\bar\theta_N}{2}},\bar\theta_N\right),$$

$$W^+:\ \ \mathcal{R}(Z_1...Z_N)=\prod_{i=1}^{N}\frac{1}{(1+\epsilon\theta_i)^{\ell_i+b_i}}\mathcal{R}\left(\frac{z_1}{1+\frac{\epsilon\theta_1}{2}},\theta_1,\frac{\bar\theta_1-\epsilon z_1}{1+\frac{\epsilon\theta_1}{2}}...\frac{z_N}{1+\frac{\epsilon\theta_N}{2}},\theta_N,\frac{\bar\theta_N-\epsilon z_N}{1+\frac{\epsilon\theta_N}{2}}\right).$$

Solutions of V^-- and W^--equations are shift-invariant functions which depend on the superdifferences only

$$\mathcal{R}(\vec{Z})=R(Z_{ij},\theta_{ij},\bar\theta_{ij}),$$

$$Z_{ij}\equiv z_i-z_j+\frac{1}{2}(\bar\theta_i\theta_j+\theta_i\bar\theta_j)\ ;\ \theta_{ij}\equiv\theta_i-\theta_j\ ;\ \bar\theta_{ij}\equiv\bar\theta_i-\bar\theta_j.$$

Let us fix some point k. There are $N-1$ independent even differences $Z_{ik}, i \neq k$ for fixed k and $2(N-1)$ independent odd differences $\theta_{ik}, \bar{\theta}_{ik}$, $i \neq k$ that transform with respect of the remaining transformations as follows

$$V^+: \quad Z_{ij} \to \frac{Z_{ij}}{(1+\frac{\epsilon\bar{\theta}_i}{2})(1+\frac{\epsilon\bar{\theta}_j}{2})} \; ; \; \theta_{ij} \to \frac{\theta_{ij} - \epsilon Z_{ij}}{(1+\frac{\epsilon\bar{\theta}_i}{2})(1+\frac{\epsilon\bar{\theta}_j}{2})} \; ; \; \bar{\theta}_{ij} \to \bar{\theta}_{ij}$$

$$W^+: \quad Z_{ij} \to \frac{Z_{ij}}{(1+\frac{\epsilon\theta_i}{2})(1+\frac{\epsilon\theta_j}{2})} \; ; \; \bar{\theta}_{ij} \to \frac{\bar{\theta}_{ij} - \epsilon Z_{ij}}{(1+\frac{\epsilon\theta_i}{2})(1+\frac{\epsilon\theta_j}{2})} \; ; \; \theta_{ij} \to \theta_{ij}.$$

It is useful to extract the representation dependent part from the function $\mathcal{R}(\vec{Z})$

$$\mathcal{R}(\vec{Z}) = \prod_{i<j} Z_{ij}^{a_{ij}} \left(1 + \frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}}\right)^{b_{ij}} R(\vec{Z}).$$

Using the formulae

$$V^+: \quad \frac{\theta_{ij}}{Z_{ij}} \to \frac{\theta_{ij}}{Z_{ij}} - \epsilon \; ; \; 1 + \frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}} \to \left(1 + \frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}}\right)(1 - \epsilon\bar{\theta}_{ij})$$

$$W^+: \quad \frac{\bar{\theta}_{ij}}{Z_{ij}} \to \frac{\bar{\theta}_{ij}}{Z_{ij}} - \epsilon \; ; \; 1 + \frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}} \to \left(1 + \frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}}\right)(1 + \epsilon\theta_{ij})$$

it is easily to check that under the conditions

$$\sum_{i\neq j} a_{ij} = -2\ell_j \; , \; a_{ij} = a_{ji} \; ; \; \sum_{i\neq j} b_{ij} = -b_j \; , \; b_{ij} = -b_{ji}$$

the equations for the function $R(\vec{Z})$ reads as follows

$$V^+: \quad R(Z_1 ... Z_N) = R\left(\frac{z_1}{1+\frac{\epsilon\bar{\theta}_1}{2}}, \frac{\theta_1 - \epsilon z_1}{1+\frac{\epsilon\bar{\theta}_1}{2}}, \bar{\theta}_1 ... \frac{z_N}{1+\frac{\epsilon\bar{\theta}_N}{2}}, \frac{\theta_N - \epsilon z_N}{1+\frac{\epsilon\bar{\theta}_N}{2}}, \bar{\theta}_N\right),$$

$$W^+: \quad R(Z_1 ... Z_N) = R\left(\frac{z_1}{1+\frac{\epsilon\theta_1}{2}}, \theta_1, \frac{\bar{\theta}_1 - \epsilon z_1}{1+\frac{\epsilon\theta_1}{2}} ... \frac{z_N}{1+\frac{\epsilon\theta_N}{2}}, \theta_N, \frac{\bar{\theta}_N - \epsilon z_N}{1+\frac{\epsilon\theta_N}{2}}\right),$$

i.e. the function $R(\vec{Z})$ is the general function of all possible superconformal invariants. Let us go to the construction of the superconformal invariants. For this purpose we fix the two points Z_j, Z_k and choose the new variables (j and k are fixed)

$$Z_{ij}^k \equiv \frac{Z_{ji}}{Z_{ik} Z_{jk}} \; , \; \theta_{ij}^k \equiv \frac{\theta_{ik}}{Z_{ik}} - \frac{\theta_{jk}}{Z_{jk}} \; , \; \bar{\theta}_{ij}^k \equiv \frac{\bar{\theta}_{ik}}{Z_{ik}} - \frac{\bar{\theta}_{jk}}{Z_{jk}}$$

with the simple transformation properties

$$V^+: \quad Z_{ij}^k \to Z_{ij}^k \cdot (1 + \epsilon\bar{\theta}_k) \; ; \; \theta_{ij}^k \to \theta_{ij}^k \; ; \; \bar{\theta}_{ij}^k \to \bar{\theta}_{ij}^k \cdot (1 + \epsilon\bar{\theta}_k)$$

$$W^+: \quad Z_{ij}^k \to Z_{ij}^k \cdot (1 + \epsilon\theta_k) \; ; \; \theta_{ij}^k \to \theta_{ij}^k \cdot (1 + \epsilon\theta_k) \; ; \; \bar{\theta}_{ij}^k \to \bar{\theta}_{ij}^k.$$

It is easy to construct the superconformal invariants using these variables. Let us fix the three points Z_j, Z_k, Z_p. Then the set of invariants is the following:

$$\frac{Z_{ij}^k}{Z_{pj}^k} = \frac{Z_{ij} Z_{pk}}{Z_{ik} Z_{pj}} \; ; \; \frac{\theta_{ij}^k \bar{\theta}_{nm}^k}{Z_{pj}^k}$$

Finally we obtain the general solution of conformal constraints

$$\mathcal{R}(\vec{z})=\prod_{i<j} z_{ij}^{a_{ij}}\left(1+\frac{\theta_{ij}\bar{\theta}_{ij}}{Z_{ij}}\right)^{b_{ij}} R\left(\frac{Z_{ij}Z_{pk}}{Z_{ik}Z_{pj}};\frac{\theta^k_{ij}\bar{\theta}^k_{nm}}{Z^k_{pj}}\right) \tag{7.0.18}$$

$$\sum_{i\neq j} a_{ij}=-2\ell_j\ ,\ a_{ij}=a_{ji}\ ;\ \sum_{i\neq j} b_{ij}=-b_j\ ,\ b_{ij}=-b_{ji}$$

where j,k,p are fixed. In the case of four-point function we can choose the superconformal invariant ansatz in the form

$$\mathcal{R}(\vec{Z})=Z_{12}^a Z_{14}^b Z_{32}^c Z_{34}^d\cdot\left(1+\frac{\theta_{12}\bar{\theta}_{12}}{Z_{12}}\right)^{\alpha}\left(1+\frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)^{\beta}\left(1+\frac{\theta_{32}\bar{\theta}_{32}}{Z_{32}}\right)^{\gamma}\left(1+\frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)^{\delta}\cdot$$

$$\cdot R\left(\frac{Z_{12}Z_{34}}{Z_{13}Z_{24}};\frac{\theta^3_{12}\bar{\theta}^3_{12}}{Z^3_{12}},\frac{\theta^3_{14}\bar{\theta}^3_{14}}{Z^3_{14}},\frac{\theta^3_{12}\bar{\theta}^3_{14}}{Z^3_{14}},\frac{\theta^3_{14}\bar{\theta}^3_{12}}{Z^3_{12}}\right)$$

where

$$a\equiv \mathrm{u}-\ell_1-\ell_2\ ;\ b\equiv -\mathrm{u}-\ell_1+\ell_2\ ;\ c\equiv -\mathrm{u}+\ell_1-\ell_2\ ;\ d\equiv \mathrm{u}+\ell_1+\ell_2$$

$$\beta=b_1-\alpha\ ;\ \gamma=-b_2-\alpha\ ;\ \delta=b_2-b_1+\alpha.$$

Appendix B

In this Appendix we collect all needed integration formulae.

General formulae of integration

$$\int_{z_4}^{z_2}\mathrm{d}Z_3 Z_{23}^A Z_{34}^B\left(1+\frac{\theta_{23}\bar{\theta}_{23}}{Z_{23}}\right)^a\left(1+\frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)^b=-\frac{\Gamma(A)\Gamma(B)}{\Gamma(A+B)}\left[\frac{Ab+Ba}{A+B}+\left(ab+\frac{AB}{4}\right)\frac{\theta_{24}\bar{\theta}_{24}}{Z_{24}}\right]Z_{24}^{A+B}$$

$$\int_{z_4}^{z_2}\mathrm{d}Z_3 Z_{23}^A Z_{34}^B\cdot\begin{pmatrix}\theta_{34}\\ \bar{\theta}_{34}\end{pmatrix}=\frac{1}{2}\frac{\Gamma(A+1)\Gamma(B+1)}{\Gamma(A+B+1)}Z_{24}^{A+B}\cdot\begin{pmatrix}\theta_{24}\\ -\bar{\theta}_{24}\end{pmatrix}$$

$$\int_{z_4}^{z_2}\mathrm{d}Z_3 Z_{23}^A Z_{34}^B\cdot\begin{pmatrix}\theta_{23}\\ \bar{\theta}_{23}\end{pmatrix}=-\frac{1}{2}\frac{\Gamma(A+1)\Gamma(B+1)}{\Gamma(A+B+1)}Z_{24}^{A+B}\cdot\begin{pmatrix}\theta_{24}\\ -\bar{\theta}_{24}\end{pmatrix}$$

Formulae of integration for even vectors

- $$\int_{z_1}^{z_3}\mathrm{d}Z_4 Z_{34}^d Z_{41}^b\left(1+\frac{\theta_{14}\bar{\theta}_{14}}{Z_{14}}\right)^{\beta}\left(1+\frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)^{\delta}\left\{1+B\frac{\theta^3_{14}\bar{\theta}^3_{14}}{Z^3_{14}}\right\}Z_{34}^n\left(1+J\frac{\theta_{34}\bar{\theta}_{34}}{Z_{34}}\right)=$$

$$=\frac{\Gamma(b)\Gamma(d+n)}{\Gamma(b+d+n)}Z_{31}^{b+d+n}\left\{C_1+C_2\frac{\theta_{31}\bar{\theta}_{31}}{Z_{31}}\right\}$$

$$C_1=B+\frac{\beta(d+n)-b(\delta+J)}{b+d+n}\ ;\ C_2=\beta(\delta+J)-\frac{b(d+n)}{4}+B\left(\delta-\beta+J+\frac{\beta(d+n)-b(\delta+J)}{b+d+n}\right)$$

- $$Z_{21}^a\left(1+\frac{\theta_{12}\bar\theta_{12}}{Z_{12}}\right)^\alpha\int_{z_1}^{z_2}\mathrm{d}Z_3Z_{23}^c\left(1+\frac{\theta_{32}\bar\theta_{32}}{Z_{32}}\right)^\gamma\left\{1+A\frac{\theta_{14}^3\bar\theta_{14}^3}{Z_{14}^3}\right\}Z_{31}^{b+d+n}\left(1+J\frac{\theta_{31}\bar\theta_{31}}{Z_{31}}\right)=$$
$$=-\frac{\Gamma(c)\Gamma(b+d+n)}{\Gamma(c+b+d+n)}Z_{21}^{a+b+c+d+n}\left\{C_1+C_2\frac{\theta_{21}\bar\theta_{21}}{Z_{21}}\right\}$$
$$C_1=A+\frac{cJ-\gamma(b+d+n)}{b+c+d+n},$$
$$C_2=-\gamma J+\frac{c(b+d+n)}{4}-\alpha\frac{cJ-\gamma(b+d+n)}{b+c+d+n}+A\left(J-\gamma-\alpha-\frac{cJ-\gamma(b+d+n)}{b+c+d+n}\right)$$

- $$Z_{21}^a\left(1+\frac{\theta_{12}\bar\theta_{12}}{Z_{12}}\right)^\alpha\int_{z_1}^{z_2}\mathrm{d}Z_3Z_{23}^c\left(1+\frac{\theta_{32}\bar\theta_{32}}{Z_{32}}\right)^\gamma\cdot$$
$$\cdot\int_{z_1}^{z_3}\mathrm{d}Z_4Z_{34}^dZ_{41}^b\left(1+\frac{\theta_{14}\bar\theta_{14}}{Z_{14}}\right)^\beta\left(1+\frac{\theta_{34}\bar\theta_{34}}{Z_{34}}\right)^\delta\frac{\theta_{14}^3\bar\theta_{12}^3}{Z_{14}^3}Z_{34}^n\left(1+J\frac{\theta_{34}\bar\theta_{34}}{Z_{34}}\right)=$$
$$=-\frac{\Gamma(b)\Gamma(d+n)}{\Gamma(b+d+n)}\frac{\Gamma(c)\Gamma(b+d+n)}{\Gamma(c+b+d+n)}Z_{21}^{a+b+c+d+n}C_1\cdot\left\{1+C_2\frac{\theta_{21}\bar\theta_{21}}{Z_{21}}\right\}$$
$$C_1=\delta+\beta+J-\frac{b+d+n}{2}\ ;\ C_2=\frac{b+d+n}{2}-\alpha-\gamma$$

Formulae of integration for odd vectors

- $$\int_{z_1}^{z_3}\mathrm{d}Z_4Z_{34}^dZ_{41}^b\left(1+\frac{\theta_{14}\bar\theta_{14}}{Z_{14}}\right)^\beta\left(1+\frac{\theta_{34}\bar\theta_{34}}{Z_{34}}\right)^\delta\left\{1+B\frac{\theta_{14}^3\bar\theta_{14}^3}{Z_{14}^3}\right\}Z_{34}^n\cdot\begin{pmatrix}\theta_{34}\\ \bar\theta_{34}\end{pmatrix}=$$
$$=\frac{\Gamma(b)\Gamma(d+n+1)}{\Gamma(b+d+n+1)}Z_{31}^{b+d+n}\cdot\begin{pmatrix}\left(\beta-\frac{b}{2}+B\right)\theta_{31}\\ \left(\beta+\frac{b}{2}+B\right)\bar\theta_{31}\end{pmatrix}$$

- $$Z_{21}^a\left(1+\frac{\theta_{12}\bar\theta_{12}}{Z_{12}}\right)^\alpha\int_{z_1}^{z_2}\mathrm{d}Z_3Z_{23}^c\left(1+\frac{\theta_{32}\bar\theta_{32}}{Z_{32}}\right)^\gamma\left\{1+A\frac{\theta_{14}^3\bar\theta_{14}^3}{Z_{14}^3}\right\}Z_{31}^{b+d+n}\cdot\begin{pmatrix}\theta_{31}\\ \bar\theta_{31}\end{pmatrix}=$$
$$=\frac{\Gamma(c)\Gamma(b+d+n+1)}{\Gamma(c+b+d+n+1)}Z_{21}^{a+b+c+d+n}\cdot\begin{pmatrix}\left(\gamma+\frac{c}{2}-A\right)\theta_{21}\\ \left(\gamma-\frac{c}{2}-A\right)\bar\theta_{21}\end{pmatrix}$$

- $$Z_{21}^a\left(1+\frac{\theta_{12}\bar\theta_{12}}{Z_{12}}\right)^\alpha\int_{z_1}^{z_2}\mathrm{d}Z_3Z_{23}^c\left(1+\frac{\theta_{32}\bar\theta_{32}}{Z_{32}}\right)^\gamma\cdot$$
$$\cdot\int_{z_1}^{z_3}\mathrm{d}Z_4Z_{34}^dZ_{41}^b\left(1+\frac{\theta_{14}\bar\theta_{14}}{Z_{14}}\right)^\beta\left(1+\frac{\theta_{34}\bar\theta_{34}}{Z_{34}}\right)^\delta\frac{\theta_{14}^3\bar\theta_{12}^3}{Z_{14}^3}Z_{34}^n\cdot\begin{pmatrix}\theta_{34}\\ \bar\theta_{34}\end{pmatrix}=$$
$$=\frac{\Gamma(b)\Gamma(d+n+1)}{\Gamma(b+d+n+1)}\frac{\Gamma(c)\Gamma(b+d+n+1)}{\Gamma(c+b+d+n+1)}Z_{21}^{a+b+c+d+n}\cdot\begin{pmatrix}0\\ \left(\gamma-\beta-\frac{b+c}{2}\right)\bar\theta_{21}\end{pmatrix}$$

References.

[1] N.M.Bogoliubov, A.G.Izergin, V.E.Korepin *Quantum inverse scattering method, correlation functions and algebraic Bethe ansatz,*
Cambridge Univ.Press 1992

[2] Z.Maassarani,J.Phys.A: **28** (1995) 1305

[3] P.B. Ramos, M.J. Martins, Nucl.Phys.**B474**, (1996) 678

[4] F.Eßler, V.Korepin , Phys. Rev. **B 46**, (1992) 9147

[5] A.Foerster, M.Karowski , Nucl.Phys. **B 396**, (1993) 611

[6] M.P.Pfannmüller and H.Frahm , Nucl.Phys. **B 479**, (1996) 575

[7] H.Frahm, M.P.Pfannmüller and A.M.Tsvelik , Phys. Rev. Lett. **81**, (1998) 2116

[8] A.Foerster, M.Karowski , Nucl.Phys. **B 408**, (1993) 512, A.Foerster, J.Phys. **A 29** (1996) 7625
D.Arnaudon JHEP **12** (1997) 006, J.Links, A.Foerster, J.Phys. **A 32** (1999) 147
A.Foerster,J.Links,A.P.Tonel, Nucl.Phys. **B 552**, (1999) 707
J.Ambjorn, D.Karakhanyan, M.Mirumyan, A. Sedrakyan *Fermionisation of the Spin-S Uimin-Lai-Sutherland Model: Generalisation of Supersymmetric t-J Model to Spin-S* cond-mat/9909432, Nucl.Phys. B[FS](2001) in press

[9] S.Derkachov, D.Karakhanyan, R.Kirschner *Heisenberg spin chains based on* $s\ell(2|1)$ *symmetry,* Nucl.Phys. **B 583**, (2000) 691

[10] E.K. Sklyanin *Classical limits of the Yang-Baxter equation,* J.Soviet.Math. **40**, (1988) 93

[11] S.E.Derkachov, V.B.Kuznetsov, E.K.Sklyanin *Baxter's Q-operator for the generic integrable XXX magnetic chain,* in preparation

[12] M.Scheunert, W.Nahm and V.Rittenberg , J.Math.Phys.**18** (1977) 155

[13] P.D.Jarvis, H.S.Green , J.Math.Phys.**20** (1979) 2115

[14] M.Marcu, J.Math.Phys.**21** (1980) 1277 , J.Math.Phys.**21** (1980) 1284

[15] D.Arnaudon, C.Chryssomalakos, L.Frappat, J.Math.Phys.**36** (1995) 5262
L. Frappat, P. Sorba, A. Sciarrino, *DICTIONARY ON LIE SUPERALGEBRAS,*
hep-th/9607161

[16] L.N. Lipatov, *High-energy asymptotics of multicolor QCD and exactly solvable lattice models,* Padova preprint DFPD-93-TH-70B, JETP.Lett. **59**(1994)596.

[17] L.D.Faddeev and G.P.Korchemsky, Phys.Lett.**B342**(1995)311.

[18] G.P.Korchemsky, Nucl. Phys. **B443**(1995)255

[19] V.Braun, S.Derkachov and A.Manashov, Phys.Rev.Lett. **81**, (1998) 2020 V.Braun, S.Derkachov, G.Korchemsky and A.Manashov, Nucl.Phys. **B 553**, (1999) 355

[20] S.Derkachov, G.Korchemsky and A.Manashov, *Evolution equations for quark-gluon distributions in multi-color QCD and open spin chains,* hep-ph/9909539, Nucl. Phys. **B566** (2000) 203.

[21] A.Belitsky, hep-ph/9907420,hep-ph/9903512, Phys.Lett. **B 453**, (1999) 59

[22] D.Karakhanian and R.Kirschner, *Conserved currents of the three-reggeon interaction,* hep-th/9902147; *High-energy scattering in gauge theories and integrable spin chains,* hep-th/9902031, Fortschr. Phys.**48**, (2000) 139

[23] H.Frahm, *Doped Heisenberg chains: spin S generalizations of the supersymmetric t-J model ,* cond-mat/9904157, Nucl.Phys.**B 559**, (1999) 613

[24] P.P. Kulish, N.Yu.Reshetikhin and E.K.Sklyanin, Lett.Math.Phys. **5** (1981) 393-403 , E.K.Sklyanin,*"Quantum Inverse Scattering Method.Selected Topics"*, in "Quantum Group and Quantum Integrable Systems" (Nankai Lectures in Mathematical Physics), ed. Mo-Lin Ge,Singapore:World Scientific,1992,pp.63-97; hep-th/9211111

[25] P.P. Kulish and E.K.Sklyanin , Zap.Nauchn.Sem. LOMI **95** (1980) 129

[26] P.P. Kulish , Zap.Nauchn.Sem. LOMI **145** (1985) 140 , J.Soviet. Math. **35** (1986) 1111, *Yang-Baxter equation and reflection equations in integrable models,*, Schladming lecture 1995, hep-th/9507070

[27] V.O.Tarasov,L.A.Takhtajan and L.D.Faddeev Theor.Math.Phys.57(1983) 163-181 ,

[28] L.D. Faddeev, Les-Houches lectures 1995, hep-th/9605187